



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-TNF-RII (p75/p80), CD120b, clone MR2-1 (Monoclonal)

Clone no. MR2-1

MONOSAN

---

|                           |  |
|---------------------------|--|
| Product name              | Mouse anti-TNF-RII (p75/p80), CD120b, clone MR2-1 (Monoclonal) |
| Host                      | Mouse  |
| Applications              | IHC-fr,FC,FUNC,ELISA,IF,IP,WB                                  |
| Species reactivity        | human, cynomolgus monkey, rhesus monkey                        |
| Conjugate                 | -  |
| Immunogen                 | Unknown or proprietary to MONOSAN and/or its suppliers         |
| Isotype                   | IgG1   |
| Clonality                 | Monoclonal   |
| Clone number              | MR2-1  |
| Size                      | 1 ml   |
| Concentration             | 100 ug/ ml   |
| Format                    | 0.2 um filtered protein G purified antibody                    |
| Storage buffer            | PBS with 0.1% BSA and 0.02% sodium azide                       |
| Storage until expiry date | 2-8°C  |

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-TNF-RII (p75/p80), CD120b, clone MR2-1 (Monoclonal)

Clone no. MR2-1

MONOSAN

**Additional info**

The antibody MR2-1 reacts with the extra-cellular part of the TNF-RII. It also reacts with the soluble receptor. TNF-RII is present on most cell types and is considered to play a prominent role in cell stimulation by TNF-alpha. TNF-RII molecule is shown to be responsible for stimulation of activated T-lymphocytes by TNF-alpha. The antibody cross reacts with rhesus and cynomolgus natural TNF-RII.

**References**

1. Leeuwenberg; JFM et al. J Immunol 1994; 152: 4036
2. Leeuwenberg, JFM et al J Immunol 1994, 152: 5070
3. Marchetti; L et al. J Biol Chem 2004; 279: 32869
4. -
5. -

**FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES**