



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Mouse anti-Citrullinated Fibrinogen Immunoglobulin, clone 3D1 (Monoclonal)

Clone no. 3D1

MONOSAN

---

Product name	Mouse anti-Citrullinated Fibrinogen Immunoglobulin, clone 3D1 (Monoclonal)
Host	Mouse
Applications	IHC(1:100-1:500), WB (1:100-1:500), ELISA(1:2000-1:4000)
Species reactivity	mouse, human
Conjugate	-
Immunogen	Deiminated murine fibrinogen peptide
Isotype	IgG1
Clonality	Monoclonal
Clone number	3D1
Size	100 ug
Concentration	n/a
Format	-
Storage buffer	lyophilized in a 10 mM ammonium bicarbonate buffer. Each vial contains 2 mg BSA
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

## Mouse anti-Citrullinated Fibrinogen Immunoglobulin, clone 3D1 (Monoclonal)

Clone no. 3D1

MONOSAN

**Additional info**

License is required for use outside research field. Fibrinogen is a protein produced by the liver which helps stop bleeding by helping blood clots to form. Fibrinogen gets deiminated (conversion from arginin to citrullin) by Peptidyl Arginine Deiminase (PAD) in inflamed joints in patients that develop rheumatoid arthritis. Citrulline, while being an amino acid, is not built into proteins during protein synthesis, as it is not coded for by DNA, yet several proteins are known to contain citrulline. Proteins that normally contain citrulline residues include myelin basic protein (MBP), filaggrin, and several histone proteins, while other proteins, like fibrin and vimentin can get deiminated during cell death and tissue inflammation. Patients with rheumatoid arthritis often (at least 80% of them) develop an immune response against proteins containing citrulline. Although the origin of this immune response is not known, detection of antibodies reactive with citrulline containing proteins or peptides is now becoming an important help in the diagnosis of rheumatoid arthritis.

References	1.	-
	2.	-
	3.	-
	4.	-
	5.	-

**FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES**