



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-Adiponectin, clone A-492 (monoclonal)

Clone no. A-492

MONOSAN

Product name	Mouse anti-Adiponectin, clone A-492 (monoclonal)
Host	Mouse
Applications	ELISA, IHC-fr, IHC-P
Species reactivity	human
Conjugate	-
Immunogen	adiponectin
Isotype	IgG1-K
Clonality	Monoclonal
Clone number	A-492
Size	100 ug
Concentration	100 ug/ml
Format	-
Storage buffer	PBS with 0.02% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Adiponectin, clone A-492 (monoclonal)

Clone no. A-492

MONOSAN

Additional info

A-492 reacts with adiponectin, an adipocytokine; adipocytokines are hormones produced in adipose tissue. Adiponectin is abundantly present in plasma and has an insulin like effect on glucose levels in the blood. Plasma adiponectin levels are found in insulin resistant patients who are obese, have diabetes mellitus type 2 or HIV-lipodystrophy. In women adiponectin levels tend to be higher than in men, which may be due to androgens suppressing adiponectin levels. Furthermore adiponectin and leptin are both indicated in regulating body weight through direct action on the hypothalamus, influencing appetite. Obese people have low adiponectin levels while levels in anorexia patients are high. Adiponectin acts as ligand for various receptors, two of which have been identified, one probably involved in carbohydrate assimilation, the other in tuning the rate of metabolism.

References

1. Dos Santos, E. et al, Oncol. Rep. 20: 971-977 (2008)
2. -
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES