



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-Fibronectin, clone 568 (monoclonal)

Clone no. 568

MONOSAN

---

|                           |  |
|---------------------------|--|
| Product name              | Mouse anti-Fibronectin, clone 568 (monoclonal)                             |
| Host                      | Mouse  |
| Applications              | FC, IHC-fr, IHC-P, IF, WB  |
| Species reactivity        | human, mouse   |
| Conjugate                 | -  |
| Immunogen                 | high molecular fraction of proteins secreted by cultured human fibroblasts |
| Isotype                   | IgG1-K   |
| Clonality                 | Monoclonal   |
| Clone number              | 568  |
| Size                      | 100 ug   |
| Concentration             | 100 ug/ml  |
| Format                    | -  |
| Storage buffer            | PBS with 0.02% sodium azide  |
| Storage until expiry date | 2-8°C  |

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

## Mouse anti-Fibronectin, clone 568 (monoclonal)

Clone no. 568

MONOSAN

**Additional info**

568 Reacts with the 8th type III repeat in the cell-binding domain of human fibronectin (220 kDa monomer; 440 kDa, dimer). Fibronectins are present in basement membranes, intestinal connective tissue matrix, and blood. Cellular fibronectin is widely distributed in the stroma of many malignant tumors.

**References**

1. Christensen, L. et al, APMIS 98(7): 615-623 (1990)
2. Christensen, L. et al, APMIS suppl. 26: 1-39 (1992)
3. Yong, J.L. et al, Int J Clin Exp Pathol 3(2): 210-216 (2010).
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES