



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-Placental Alkaline Phosphatase (PLAP), clone F5C2 (C2) (Monoclonal)

Clone no. F5C2

MONOSAN

Product name	Mouse anti-Placental Alkaline Phosphatase (PLAP), clone F5C2 (C2) (Monoclonal)
Host	Mouse
Applications	ELISA, FC, IHC-fr, IHC-P, IP, WB
Species reactivity	human
Conjugate	-
Immunogen	PLAP, purified from FS phenotype human placenta
Isotype	IgG1
Clonality	Monoclonal
Clone number	F5C2
Size	100 ug
Concentration	100 ug/ml
Format	-
Storage buffer	PBS with 0.02% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Placental Alkaline Phosphatase (PLAP), clone F5C2 (C2) (Monoclonal)

Clone no. F5C2

MONOSAN

Additional info

PLAP is a tissue specific, throphoblast-derived, 58 kDa, glycosyl-phosphatidylinositol (GPI)- anchored, dimeric, Zn²⁺ metallated glycoprotein, only found in humans, orangutans and chimpanzees, that catalyzes the hydrolysis of phosphomonoesters into an inorganic phosphate and an alcohol. It is present in the placenta and serum of pregnant women and in high frequency in gynecological and testicular cancers and in lower frequency in other tumors. The three tissue-specific APs in humans, PLAP, germ cell AP (GCAP) and intestinal AP, are 90-98% homologous. Non tissue specific AP is found in kidney, liver and bone. F5C2 binds equally well to all common allelic variants (S,F, FS and I) of PLAP and to some variants of AP from normal human testis, while antibody H7E8 reacts with all variants of AP in normal human testis.

References

1. Millan J.L. et al, Eur. J. Biochem. 136: 1-7 (1983)
2. -
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES