



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-CD43, clone 84-3C1 (Monoclonal)

Clone no. 84-3C1

MONOSAN

Product name	Mouse anti-CD43, clone 84-3C1 (Monoclonal)
Host	Mouse
Applications	FC, IHC-fr, IHC-P, IF, WB
Species reactivity	human
Conjugate	-
Immunogen	stimulated human leucocytes
Isotype	IgG1-K
Clonality	Monoclonal
Clone number	84-3C1
Size	100 ug
Concentration	100 ug/ml
Format	-
Storage buffer	PBS with 0.02% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-CD43, clone 84-3C1 (Monoclonal)

Clone no. 84-3C1

MONOSAN

Additional info

84-3C1 reacts with a 95/115 kDa protein on T-cells and thymocytes and a 115/135 kDa molecule on neutrophils and platelets. 70-90% of T-cell lymphomas and from 22-37% of Bcell lymphomas express CD43. No reactivity has been observed with reactive B-cells. So a Blineage population that co-expresses CD43 is highly likely to be a malignant lymphoma, especially a low-grade lymphoma, rather than a reactive B-cell population. When CD43 antibody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD43 argues against a reactive process and favors a diagnosis of lymphoma. In addition, expression is altered in Wiskott Aldrich syndrome. A proportion of AIDS patients have antibodies to CD43. A soluble form called galactoglycoprotein is present in serum. 85-3C1 was typed at the 3rd International Workshop on Human Leucocyte Differentiation antigens.

References

1. Cobbold, S. et al. In Leucocyte typing III, Oxford University Press, pp 789-803 (1989)
2. Stross, WP. et al. J. Clin. Path. 42: 953-961 (1989)
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES