



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-CD80, clone C80-99 (Monoclonal)

Clone no. C80-99

MONOSAN

---

|                           |  |
|---------------------------|--|
| Product name              | Mouse anti-CD80, clone C80-99 (Monoclonal) |
| Host                      | Mouse                                      |
| Applications              | FC, IHC-fr, IF                             |
| Species reactivity        | human                                      |
| Conjugate                 | -  |
| Immunogen                 | recombinant full length CD80               |
| Isotype                   | IgG2b-K                                    |
| Clonality                 | Monoclonal                                 |
| Clone number              | C80-99                                     |
| Size                      | 100 ug                                     |
| Concentration             | 100 ug/ml                                  |
| Format                    | -  |
| Storage buffer            | PBS with 0.02% sodium azide                |
| Storage until expiry date | 2-8°C                                      |

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-CD80, clone C80-99 (Monoclonal)

Clone no. C80-99

MONOSAN

**Additional info**

CD80 is involved in the costimulatory signal essential for T-lymphocyte activation. T-cell proliferation and cytokine production is included by the binding of CD28, binding to CTLA-4 has opposite effects and inhibits T-cell activation.

**References**

1. Peach, R.J., et al. J. Biol. Chem. 270: 21181-21187 (1995)
2. Fargeas, C.A., et al. J. Exp. Med. 182: 667-675 (1995)
3. -
4. -
5. -

**FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES**