



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-Cyclin D1, clone P2D11F11 (monoclonal)

Clone no. P2D11F11

MONXtra

---

|                           |   |
|---------------------------|---|
| Product name              | Mouse anti-Cyclin D1, clone P2D11F11 (monoclonal)                         |
| Host                      | Mouse   |
| Applications              | IHC-P (1:50)  |
| Species reactivity        | human   |
| Conjugate                 | -   |
| Immunogen                 | Prokaryotic fusion protein corresponding to the human cyclin D1 molecule. |
| Isotype                   | IgG2a   |
| Clonality                 | Monoclonal  |
| Clone number              | P2D11F11  |
| Size                      | 1 ml  |
| Concentration             | Greater than or equal to 19 mg/L  |
| Format                    | -   |
| Storage buffer            | Tissue culture supernatant with Sodium azide                              |
| Storage until expiry date | 2-8°C   |

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Cyclin D1, clone P2D11F11 (monoclonal)

Clone no. P2D11F11

MONXtra

---

**Additional info**

The D-type cyclins are a family of proteins which function primarily by regulating the activity of cyclin dependent kinases in the G1 phase of the cell cycle. Cyclin D1, a protein of 36 kD, is also known as PRAD1 or bcl-1. Maximum expression of cyclin D1 occurs at a critical point in mid to late G1 phase of the cell cycle. The cyclin D1 gene, located on 11q13 has been reported to be overexpressed in mantle cell lymphomas due to the chromosomal translocation t(11;18).

**References**

1. McIntosh GG et al. *Oncogene*. 1995; 11:885–891
2. Saiz AD et al. *Journal of Pathology*. 2002; 198(2):157–162
3. Mommers ECM et al. *Journal of Pathology*. 2001; 194(3):327–333
4. Saito T et al. *Journal of Pathology*. 2001; 195(2):222–228
5. Sheyn I et al. *Human Pathology*. 1997; 28(3):270–276

**FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES**