



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

Mouse anti-CD5 (Liquid Primary Antibody), clone 4C7 (monoclonal)

Clone no. 4C7

MONXtra

---

Product name	Mouse anti-CD5 (Liquid Primary Antibody), clone 4C7 (monoclonal)
Host	Mouse
Applications	IHC-P (1:100)
Species reactivity	human
Conjugate	-
Immunogen	Recombinant prokaryotic fusion protein corresponding to the external domain of the human CD5 molecule.
Isotype	IgG1, kappa
Clonality	Monoclonal
Clone number	4C7
Size	1 ml
Concentration	Greater than or equal to 76 mg/L
Format	-
Storage buffer	Tissue culture supernatant with sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-CD5 (Liquid Primary Antibody), clone 4C7 (monoclonal)

Clone no. 4C7

MONXtra

---

**Additional info**

CD5 antigen is reported to be expressed on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed on T cells. CD5 antigen is also expressed by many T cell leukemias, lymphomas, activated T cells and on a subset of B cells located primarily in the mantle zones of normal lymph nodes. CD5 antigen expression is also reported in T cell acute lymphocytic leukemias (T-ALL), some B cell chronic lymphocytic leukemias (B-CLL) as well as B and T cell lymphomas.

**References**

1. Leong FJW et al. The Journal of Histotechnology. 2002; 25(4):215-227
2. Walsh R et al. Arch. of Pathol.and Lab.Medicine. 2001; 125(6):781-784
3. Tateyama H et al. American Journal of Clinical Pathology. 1999; 111(2):235-240
4. Dorfman DM & Shamsafaei A. Modern Pathology. 1997; 10(9):859-863
5. Kaufmann O et al. American Journal of Clinical Pathology. 1997; 108(6):669-673

**FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES**