



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-CD34 Endothelial Cell Marker, clone QBEnd/10 (monoclonal)

Clone no. QBEnd/10

MONXtra

---

|                           |  |
|---------------------------|--|
| Product name              | Mouse anti-CD34 Endothelial Cell Marker, clone QBEnd/10 (monoclonal)                         |
| Host                      | Mouse  |
| Applications              | IHC-P (1:100)  |
| Species reactivity        | human  |
| Conjugate                 | -  |
| Immunogen                 | Detergent solubilized vesicular suspension prepared from a perfusate of human term placenta. |
| Isotype                   | IgG1   |
| Clonality                 | Monoclonal   |
| Clone number              | QBEnd/10   |
| Size                      | 1 ml   |
| Concentration             | Greater than or equal to 261 mg/L  |
| Format                    | -  |
| Storage buffer            | Tissue culture supernatant with Sodium azide   |
| Storage until expiry date | 2-8°C  |

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-CD34 Endothelial Cell Marker, clone QBEnd/10 (monoclonal)

Clone no. QBEnd/10

MONXtra

---

**Additional info**

The CD34 antigen is a single chain transmembrane glycoprotein with a molecular weight of 110 kD. The CD34 protein is selectively expressed on human lymphoid and myeloid hemopoietic progenitor cells. The CD34 antigen is also expressed on vascular endothelium. Enzyme digestion of paraffin sections is recommended with clone QBEnd/10 in preference to heat induced epitope retrieval as it produces stronger staining and reduces background elastin staining

**References**

1. Tornoczky T et al. Journal of Clinical Pathology. 2003; 56(5):363–367
2. Cox G et al. Lung Cancer. 2000; 29(3):169–177
3. Dhillon AP et al. Journal of Pathology. 1990; 162:274
4. Ramani P et al. Histopathology. 1990; 17(3):237–242
5. Watanabe T et al. Journal of Clinical Pathology. 2001; 54:631–636.

**FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES**