



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-Cytokeratin 5 (Liquid Primary Antibody), clone XM26 (monoclonal)

Clone no. XM26

MONXtra

Product name	Mouse anti-Cytokeratin 5 (Liquid Primary Antibody), clone XM26 (monoclonal)
Host	Mouse
Applications	IHC-P (1:100)
Species reactivity	human
Conjugate	-
Immunogen	Prokaryotic recombinant fusion protein corresponding to a 103 amino acid portion of the C-terminal region of the human cytokeratin 5 molecule.
Isotype	IgG1, kappa
Clonality	Monoclonal
Clone number	XM26
Size	1 ml
Concentration	Greater than or equal to 21 mg/L
Format	-
Storage buffer	Tissue culture supernatant with Sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Cytokeratin 5 (Liquid Primary Antibody), clone XM26 (monoclonal)

Clone no. XM26

MONXtra

Additional info

Cytokeratins are a large family of cytoskeletal proteins found in epithelial cells. They are co-ordinately synthesized in pairs so that at least one member of each family is expressed in each epithelial cell. Cytokeratins assemble into obligatory heteropolymers composed of type I (acidic) and type II (basic) polypeptides to form higher order tetramers and protofilaments. Basal cells of human epidermis express acidic keratin 14 and basic cytokeratin 5. Cytokeratin 5 is a 58 kD protein that is closely related to cytokeratin 6. Point mutations in the cytokeratin 5 gene at locus 12q11-q13 can cause various types of epidermolysis bullosa simplex. Cytokeratin 5 is also reported to be expressed in most epithelial and biphasic mesotheliomas. Clone XM26 is specific for the 58 kD intermediate filament protein known as cytokeratin 5. It is not cross-reactive with cytokeratin 6.

References

1. Bhargava R et al. The American Journal of Clinical Pathology . 2008; 130:724-731
2. Laakso M et al. Clinical Cancer Research. 2006; 12(14):4185-4191
3. Miettinen M et al. American Journal of Surgical Pathology. 2003; 27(2):150–158
4. Zhang RR et al. Breast Cancer Research. 2003; 5:R151–R156
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES