



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-Alpha-Synuclein, clone KM51 (monoclonal)

Clone no. KM51

MONXtra

| | |
|---------------------------|---|
| Product name | Mouse anti-Alpha-Synuclein, clone KM51 (monoclonal) |
| Host | Mouse |
| Applications | IHC-P (1:20-1:40) |
| Species reactivity | human |
| Conjugate | - |
| Immunogen | Prokaryotic recombinant protein corresponding to the majority of the full length alpha-synuclein molecule |
| Isotype | IgG1, kappa |
| Clonality | Monoclonal |
| Clone number | KM51 |
| Size | 1 ml |
| Concentration | n/a |
| Format | - |
| Storage buffer | Tissue culture supernatant with 15mM sodium azide |
| Storage until expiry date | 2-8°C |

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Alpha-Synuclein, clone KM51 (monoclonal)

Clone no. KM51

MONXtra

Additional info

Alpha-synuclein is a protein of 140 amino acids and a member of the synuclein family. It shares 61% sequence homology with beta-synuclein and is highly conserved between vertebrate species. It does not possess a signal sequence suggesting that it is an intracellular protein. All synucleins have an unusual organization based around the eleven residue repeating motif and an alpha-helical secondary structure resembling those found in the lipid-binding domain of exchangeable apolipoproteins, including Apo E. This homology suggests a direct interaction of alpha-synuclein with membranes consistent with its affinity for synaptosomes. Clone KM51 is specific for alpha-synuclein and is unreactive with beta-synuclein. Pretreatment of tissue sections with 98-100% formic acid is also recommended.

References

1. Baba M et al. American Journal of Pathology. 152 (4): 879–884 (1998)
2. Polymeropoulos MH et al. Science. 276: 2045–2047 (1997)
3. Spillantini MG et al. Nature. 388: 839–840 (1997)
4. Weinreb PH et al. Biochemistry. 35 (43): 13709–13715 (1996)
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES