



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-Multiple Myeloma Oncogene 1 (MUM-1), Clone EAU32 (monoclonal)

Clone no. EAU32

MONXtra

| | |
|---------------------------|---|
| Product name | Mouse anti-Multiple Myeloma Oncogene 1 (MUM-1), Clone EAU32 (monoclonal) |
| Host | Mouse |
| Applications | IHC-P (1:100) |
| Species reactivity | human |
| Conjugate | - |
| Immunogen | Prokaryotic recombinant protein corresponding to 313 amino acids of the human multiple myeloma oncogene 1 (MUM-1) molecule. |
| Isotype | IgG1 |
| Clonality | Monoclonal |
| Clone number | EAU32 |
| Size | 1 ml |
| Concentration | Greater than or equal to 263 mg/L |
| Format | - |
| Storage buffer | Tissue culture supernatant with sodium azide |
| Storage until expiry date | 2-8°C |

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Multiple Myeloma Oncogene 1 (MUM-1), Clone EAU32 (monoclonal)

Clone no. EAU32

MONXtra

Additional info

The MUM-1 (multiple myeloma oncogene 1) gene was originally identified because of its involvement in the t(6:14) translocation observed in multiple myeloma, which causes the juxtaposition of the MUM-1 gene to the Ig heavy chain locus. MUM-1 is expressed in late plasma cell directed stages of B cell differentiation and in activated T cells, suggesting that MUM-1 may serve as a marker for lympho-hemopoietic neoplasms derived from these cells. The morphologic spectrum of MUM-1 expressing cells has been found to range from that of a centrocyte to that of a plasmablast/plasma cell. Consequently the histogenic value of MUM-1 may be to provide a marker to aid in the identification of the transition from BCL-6 positive (germinal center B cells) to CD138 positive (immunoblasts and plasma cells).

References

1. Bergsagel P and Kuehl W. Oncogene. 2001; 20(40):5611-5622
2. Iida S et al. Nature Genetics. 1997;17(2):226-230
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES