



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Product datasheet PS040

**MONOSAN**<sup>®</sup>

Polyclonal Rabbit anti-Laminin  
Clone no. -

MONOSAN

---

Product name	Polyclonal Rabbit anti-Laminin
Host	Rabbit
Applications	IHC-fr, IHC-P (1:50 - 1:100), WB
Species reactivity	Bovine, Human, Mouse, Pig, Rabbit, Rat
Conjugate	-
Immunogen	Laminin isolated from Engelbrecht-Holm-Swarm (EHS)-mouse sarcoma
Isotype	-
Clonality	Polyclonal
Clone number	-
Size	250 µl
Concentration	n/a
Format	-
Storage buffer	Contains 0.09% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Polyclonal Rabbit anti-Laminin

Clone no. -

MONOSAN

---

**Additional info**

Laminin is a glycoprotein (Mr 850 - 1.000 kD, consisting of 3 glycosylated polypeptide chains with molecular weights of 440 and 225 (2x) kD) produced by various human epithelial and mesenchymal cells, and forms an extracellular matrix of thin filaments. In normal tissues, laminin is invariably present in all basal laminas surrounding muscle, nerve, fat and decidua cells and separates epithelial and endothelial cells from abutting connective tissues. Laminin has also been identified within the cytoplasm of breast epithelia, stromal cells of the endometrium, and within endothelial, bile duct epithelial and mesenchymal cells of the liver. Laminin has been found to be involved in cellular activities such as adhesion, spreading, differentiation, polarization, proliferation, locomotion, tissue invasion and chemotactic responses.

No cross reaction was obtained with human type I, III, IV and V collagen in immunoblotting, whereas the antibody reacted with a distinct band of appr. 200-220 kD from a 8M Urea extract from amnion basement membrane.

Positive control: skin, kidney.

**References**

1. Domke, L et al. Cell Tissue Res 2019;375:451-482
2. Costa, B. et al. Blood.Adv. 2019;3:1092-1102
3. Oh, J. et al. Oncogene 2004;23: 5041-5048
4. Eisemann, T. et al. BMC.Cancer 2018;18:103
5. Eisemann, T. et al. Neuro.Oncol. 2019;21: 326-336

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES