



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Product datasheet PS286

MONOSAN[®]

Polyclonal Bovine anti-Lactoferrin

Clone no. -

MONOSAN

| | |
|---------------------------|--|
| Product name | Polyclonal Bovine anti-Lactoferrin |
| Host | Bovine |
| Applications | ELISA,IP,WB |
| Species reactivity | human |
| Conjugate | - |
| Immunogen | Unknown or proprietary to MONOSAN and/or its suppliers |
| Isotype | Ig |
| Clonality | Polyclonal |
| Clone number | - |
| Size | 1 ml |
| Concentration | 100 ug/ ml |
| Format | - |
| Storage buffer | PBS with 0.1% BSA and 0.02% sodium azide |
| Storage until expiry date | 2-8°C |

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Polyclonal Bovine anti-Lactoferrin

Clone no. -

MONOSAN

Additional info

Human lactoferrin (LF) is an 80 kDa glycoprotein which was first isolated from human milk. It plays an important part in the immune system and helps to fight infections. Lactoferrin promotes the health of the gastro-intestinal system by improving the intestinal microbial balance. In addition, LF can be found in epithelia and most body fluids and secretions. Lactoferrin is secreted in plasma by neutrophils. Its plasma concentration also represents a positive relation to the total pool of neutrophils and the rate of neutrophil turnover. In inflammation lactoferrin is released from secondary granules of neutrophilic leukocytes into the extracellular medium. Therefore the extracellular lactoferrin concentration can be used as an index for neutrophil activation. Lactoferrin strongly binds to iron and this iron binding property is considered to be an important antimicrobial. Human lactoferrin binds to bacterial products through its highly positively charged N-terminus, it kills various bacteria, most probably by inducing intracellular changes in these bacteria without affecting the membrane permeability. Cleavage by pepsin of lactoferrin leads to the release of lactoferricin H. This 47 amino acid peptide has more antimicrobial activity than its precursor and it can inhibit the classical but not the alternative complement pathway. Lactoferrin also plays a role in signal transduction, immunomodulation and has antiadhesive, anticancer, antiviral activity.

References

1. Nuijens; J et al. J Biol Chem 1997; 272: 8802
2. Nibbering, P et al Infect Immun 2001, 69: 1469
3. Veen; van H et al. Anal Biochem 2002; 309: 60
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES