



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Polyclonal Antibody to Human Factor VIII Related Antigen

Description:

- Immunogen:** Factor VIII related antigen isolated from human plasma.
- Format:** This antibody has been pretitered and quality controlled to work on formalin-fixed paraffin-embedded as well as acetone fixed cryostat tissue sections. No further titration is required.
- Specificity:** This antibody reacts with factor VIII related antigen. It stains endothelial cells and megakaryocytes in various human tissues. This antibody does not cross-react with any other cell types.

Uses/Limitations: Immunohistochemistry
For Research Use Only.
Do not use past expiration date.

Storage: 2-8° Centigrade.

Procedure: We suggest an incubation period of 30-60 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Although not required, proteolytic treatment may enhance the immunostaining in overfixed tissues.

Precautions: Contains Sodium Azide as a preservative.
Do not pipette by mouth.

References:

- i) Bukh et al. Thromb Res 43: 579, 1986.
- ii) Cejka et al. Clin Chem 28: 1356, 1982.
- iii) Sehested et al. Virchows Arch (Pathol Anat) 391: 217, 1981.