



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Kappa; Clone L1C1 (Ready-To-Use)

| <b>Availability/Contents:</b> | <u>Item #</u> | <u>Volume</u> |
|-------------------------------|---------------|---------------|
|                               | A00113-0002   | 2 ml          |
|                               | A00113-0007   | 7 ml          |
|                               | A00113-0025   | 25 ml         |

**Description:**

**Species:** Mouse

**Immunogen:** B lymphoma cells.

**Clone:** L1C1

**Isotype:** Mouse IgG1, Kappa

**Format:** This antibody has been pretitered and quality controlled to work on formalin-fixed paraffin-embedded as well as acetone fixed cryostat tissue sections. No further titration is required.

**Specificity:** The kappa light chain antibody recognizes the kappa light chain of immunoglobulin.

**Background:** The kappa light chain itself is one of the two small polypeptide subunits of an antibody, the other being lambda. Antibodies are produced by the B-cells in lymphoid tissue. Each B lymphocyte expresses either lambda or kappa light chain but never both together. Hence, the kappa light chain antibody is a useful marker for identifying B lymphocytes expressing kappa light chain.

If a lymph node or other tissue of lymphoid origin is normal or benign, it should contain a mixture of lambda and kappa light chain positive cells. However, if there is only one type, such as all kappa light chain positive, then they may have all been derived from a clonal population. This may be indicative of a pathological condition, including a malignancy. As such, the kappa light chain antibody has been reported to help identify leukemias, plasmacytomas and certain non-Hodgkin's lymphomas. The underlying mechanism of identification by the kappa light chain antibody in these various cancers would be their expression of the kappa, but not lambda, light chain.

Hence, the kappa light chain antibody has an overall usefulness in identifying normal B-cells expressing kappa light chain as well as helping to identify malignancies or potentially other pathologies characterized by a clonally derived kappa light chain positive population.

**Species Reactivity:** Human.


**Positive Control:** Tonsil.

**Cellular Localization:** Cytoplasmic.

**Titer/Working Dilution:** No further dilution is required.


**Microbiological State:** This product is not sterile.

Storage: 2° C  8° C




ScyTek Laboratories, Inc.  
205 South 600 West  
Logan, UT 84321  
U.S.A.

**CE**

 EmergoEurope (31)(0) 70 345-8570  
Molsnstraat 15  
2513 BH Hague, The Netherlands

**Uses/Limitations:** Not to be taken internally.  
 For Research Use Only.  
 This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy.  
 Do not use if reagent becomes cloudy.  
 Do not use past expiration date.  
 Use caution when handling reagents.  
 Non-Sterile.



Ordering Information and Current Pricing at [www.scytek.com](http://www.scytek.com)

**Procedure:**

1. **Tissue Section Pretreatment:** Staining of formalin fixed, paraffin embedded tissue sections is enhanced by pretreatment with Citrate Plus (ScyTek catalog# CPL500) or Citrate Buffer (10x), pH 6.0 (ScyTek Catalog# CBB500, see IFU for instructions).
2. **Primary Antibody Incubation Time:** We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user.
3. **Visualization:** For maximum staining intensity we recommend the “UltraTek HRP Anti-Polyvalent Lab Pack” (ScyTek catalog# UHP125, see IFU for instructions) combined with the “DAB Chromogen/Substrate Bulk Pack (High Contrast)” (ScyTek catalog# ACV500, see IFU for instructions).

**Precautions:** Contains Sodium Azide as a preservative (0.09% w/v).  
 Do not pipette by mouth.  
 Avoid contact of reagents and specimens with skin and mucous membranes.  
 Avoid microbial contamination of reagents or increased nonspecific staining may occur.  
 This product contains no hazardous material at a reportable concentration according to U.S. 29 CFR 1910.1200, OSHA Hazardous Communication Standard and EC Directive 91/155/EC.


**References:**

1. Korkolopoulou P, GA Pangalis E Patsouris, VaA Boussiotis, C Kittas C. Leukemia Lymphoma, 1994, 13:151-159 (1994).
2. Abe M, T Goto, SJ Kennel, D Wolfenbarger, SD Macy, DT Weiss, A Solomon. AJCP 100:67-74 (1993).
3. Berg AM, RF Troxler, G Grillone, J Kasznica, K Kane, AS Cohen, M Skinner. Localized amyloidosis of the larynx: evidence for light chain composition. Ann. Otol. Rhinol. Laryngol. 884-889 (1993).
4. Takahashi H, S Fujita, H Okabe, N Tsuda, F Tezuka. Pathol Res Prac 189:300-311 (1993).
5. Momose H, YY Chen, J Ben-Ezra, LM Weiss. Hum Pathol. 23:1115-1119 (1992).


**Warranty:**

No products or “Instructions For Use (IFU)” are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our IFU or website. Our warranty is limited to the actual price paid for the product. ScyTek Laboratories, Inc. is not liable for any property damage, personal injury, time or effort or economic loss caused by our products. Immunohistochemistry is a complex technique involving both histological and immunological detection methods. Tissue processing and handling prior to immunostaining can cause inconsistent results. Variations in fixation and embedding or the inherent nature of the tissue specimen may cause variations in results. Endogenous peroxidase activity or pseudoperoxidase activity in erythrocytes and endogenous biotin may cause non-specific staining depending on detection system used.

Storage: 2° C  8° C

 ScyTek Laboratories, Inc.  
 205 South 600 West  
 Logan, UT 84321  
 U.S.A.

**CE**

 EmergoEurope (31)(0) 70 345-8570  
 Molsnstraat 15  
 2513 BH Hague, The Netherlands