



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Peroxide Block

Description: Peroxide Block has been developed to use with immunolabeling techniques for the reduction of nonspecific background staining do to endogenous peroxidase.

Availability/Contents:	Item #	Volume
	ACA015	15 ml Dropper
	ACA125	125 ml
	ACA500	500 ml
	ACA999	1000 ml

Uses/Limitations:

- Not to be taken internally.
- For In-Vitro Diagnostic use.
- Histological applications.
- Do not use if reagents become cloudy.
- Do not use past expiration date.
- Use caution when handling reagents.
- Non-Sterile.



Ordering Information and Current Pricing at www.scytek.com

Precautions:

- Avoid contact with skin and eyes.
- Harmful if swallowed.
- Follow all Federal, State, and local regulations regarding disposal.

Procedure:


1. Prior to application of primary antibody, apply Peroxide Block to tissue section and incubate slide for 10 minutes.
2. Rinse 1 time in buffer.

Note: For bulk staining, pour Peroxide Block in a covered staining tray and dip slides for 5-10 minutes. Replace with fresh reagent after 5-10 uses

References:

1. Duncan SM, Seigel GM. High-contrast enzymatic immunohistochemistry of pigmented tissues. Journal of biological methods. 2016;3(3).
2. Firat U, Kaya S, Çim A, Büyükbayram H, Gökalp O, Dal MS, Tamer MN. Increased caspase-3 immunoreactivity of erythrocytes in STZ diabetic rats. Experimental diabetes research. 2012 Apr 26;2012.

Storage: 2° C  8° C



ScyTek Laboratories, Inc.
205 South 600 West
Logan, UT 84321
U.S.A.

CE 

EC REP

Emergo Europe
Prinsessegracht 20
2514 AP The Hague, The Netherlands