



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Permanent Red Bulk Pack (For Alkaline Phosphatase)

**Description:** Permanent Red (For Alkaline Phosphatase) is a two-component chromogen for immunohistochemical staining and immunoblotting. When in the presence of alkaline phosphatase enzyme, Permanent Red produces an intense red precipitate that may be permanently mounted in synthetic resin.

**Uses/Limitations:** Not to be taken internally.  
For In-Vitro Diagnostic use.  
Histological applications.  
Do not use if reagents become cloudy.  
Do not use past expiration date.  
Use caution when handling reagents.  
Non-Sterile.



**Control Tissue:** Any well fixed paraffin embedded or frozen tissue section.

**Ordering information regarding individual components on back page!**


### Kit Options:


<u>Item #</u>	<u>Component</u>	<u>Volume</u>	<u>Storage</u>
PRD500	Permanent Red Concentrate	15 ml	2-8°C
	Permanent Red Buffer	500 ml	2-8°C
PRD999	Permanent Red Concentrate	30 ml	2-8°C
	Permanent Red Buffer	1000 ml	2-8°C

**Precautions:** Avoid contact with skin and eyes.  
Harmful if swallowed.  
Follow all Federal, State, and local regulations regarding disposal.

### Procedure:

- Combine 10µl of Permanent Red Concentrate with each 1ml of Permanent Red Buffer. Used combined mixture immediately after mixing.
- Dip slide one time in DI/Distilled water immediately prior to application of Permanent Red.
- Apply mixture to tissue section and incubate for 10 minutes.

Storage: 2° C  8° C

 ScyTek Laboratories, Inc.  
205 South 600 West  
Logan, UT 84321  
U.S.A.

CE IVD


EC REP  
Emergo Europe  
Prinsessegracht 20  
2514 AP The Hague, The Netherlands


P.O. Box 3286 - Logan, Utah 84323, U.S.A. - Tel. (800) 729-8350 – Tel. (435) 755-9848 - Fax (435) 755-0015 - [www.scytek.com](http://www.scytek.com)

4. Rinse slide using DI/Distilled water.
5. If desired, repeat combining and application of mixture.
6. Rinse slide in two changes of DI/Distilled water.
7. Counterstain as desired.
8. Quickly dehydrate in alcohol and clear in xylene or substitute.  
*Note: Alcohol and Xylene can cause chromogen to leach from tissue over extended periods of time.*
9. Mount and coverslip with synthetic medium.

Bulk Reagent Ordering Information and Current Pricing at [www.scytek.com](http://www.scytek.com)

Description:	Catalog #	Volume
Permanent Red Concentrate	PRC015	15 ml
	PRC030	30 ml
Permanent Red Buffer	PRB500	500 ml
	PRB999	1000 ml

Storage: 2° C  8° C

 ScyTek Laboratories, Inc.  
205 South 600 West  
Logan, UT 84321  
U.S.A.



Emergo Europe  
Prinsessegracht 20  
2514 AP The Hague, The Netherlands