

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





Instructions For Use

RA0134-C.5-IFU-RUO

Rev. Date: Oct. 21, 2014

Revision: 1

Page 1 of 2

P.O. Box 3286 - Logan, Utah 84323, U.S.A. - Tel. (800) 729-8350 - Tel. (435) 755-9848 - Fax (435) 755-0015 - www.scytek.com

HSP60 (Heat Shock Protein 60); Clone HLD4/780

(Concentrate)

Availability/Contents: Item # Volume
RA0134-C.5 Volume
0.5 ml

Description:

Species: Mouse

Immunogen: Recombinant human HSPD1 protein

Clone: HLD4/780 Isotype: IgG1, kappa

Entrez Gene ID: 3329 (Human); 15510 (Mouse); 63868 (Rat)

Hu Chromosome Loc.: 2q33.1

Synonyms: 60kDa chaperonin; 60kDa heat shock protein mitochondrial; Chaperonin; 60-KD (CPN60);

GROEL; HLD4; HSP65; HSPD1; HuCHA60; Mitochondrial matrix protein P1; P60 lymphocyte

protein; Short heat shock protein 60 Hsp60s1; Spastic paraplegia 13 (SPG13).

Mol. Weight of Antigen: 60kDa

Format: 200µg/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS

with 0.05% BSA & 0.05% azide.

Specificity: Recognizes a 60kDa protein, identified as the heat shock protein 60 (HSP60).

Background: A wide variety of environmental and pathophysiological stressful conditions trigger the synthesis

of a family of proteins known as heat shock proteins (HSP's), more appropriately called stress response proteins (SRP's). HSP60 is a potential antigen in a number of autoimmune diseases. In human arthritis and in experimentally induced arthritis in animals, disease development coincides with the development of immune reactivity directed against not only bacterial HSP60,

but also against its mammalian homolog.

Species Reactivity: Human, Mouse, Rat, Hamster, Sheep, Rabbit, Cow, Dog, Pig, Monkey, Chicken, Xenopus

laevis, and Drosophila. Does not react with Bacteria, Helminths, and Spinach. Others not

known.

Positive Control: HeLa or HepG2 cells. Breast carcinoma. Synovial biopsies from patients with juvenile chronic

arthritis. Synovial lining layer is strongly positive for HSP60.

Cellular Localization: Cytoplasmic

Titer/ Working Dilution: Immunohistochemistry (Frozen and Formalin-fixed): 0.5-1 μg/ml

Flow Cytometry: 0.5-1 µg/million cells

 $\begin{array}{ll} \mbox{Immunofluorescence:} & 0.5\mbox{-}1 \ \mbox{μg/ml$} \\ \mbox{Western Blotting:} & 0.25\mbox{-}0.5 \ \mbox{μg/ml$} \\ \end{array}$

Immunoprecipitation: 0.5-1 μg/500μg protein lysate

Microbiological State: This product is not sterile.

Storage: 2° C 8° C

ScyTek Laboratories, Inc. 205 South 600 West Logan, UT 84321 U.S.A.

CE

EmergoEurope (31)(0) 70 345-8570 Molsnstraat 15 2513 BH Hague, The Netherlands



Instructions For Use RA0134-C.5-IFU-RUO

Rev. Date: Oct. 21, 2014

Revision: 1

Page 2 of 2

P.O. Box 3286 - Logan, Utah 84323, U.S.A. - Tel. (800) 729-8350 - Tel. (435) 755-9848 - Fax (435) 755-0015 - www.scytek.com

Uses/Limitations: Not to be taken internally.

For Research Use Only.


This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded

tissue sections, to be viewed by light

microscopy.

Do not use if reagent becomes cloudy. Do not use past expiration date.

Non-Sterile.



Formalin-paraffin breast carcinoma stained with HSP60: Clone HLD4/780.

Ordering Information and Current Pricing at www.scytek.com

Procedure:

- 1. **Tissue Section Pretreatment (Highly Recommended):** Staining of formalin fixed, paraffin embedded tissue sections is significantly enhanced by pretreatment with Citrate Plus (ScyTek catalog# CPL500).
- Primary Antibody Incubation Time: We suggest an incubation period of 30 minutes at room temperature.
 However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user.
- 3. **Visualization:** For maximum staining intensity we recommend the "UltraTek HRP Anti-Polyvalent Lab Pack" (ScyTek catalog# UHP125, see IFU for instructions) combined with the "DAB Chromogen/Substrate Bulk Pack (High Contrast)" (ScyTek catalog# ACV500, see IFU for instructions).

Precautions:

Contains Sodium Azide as a preservative (0.09% w/v).

Do not pipette by mouth.

Avoid contact of reagents and specimens with skin and mucous membranes.

Avoid microbial contamination of reagents or increased nonspecific staining may occur.

This product contains no hazardous material at a reportable concentration according to U.S. 29 CFR 1910.1200,

OSHA Hazardous Communication Standard and EC Directive 91/155/EC.

References:

Boog CJ; de Graeff-Meeder ER; Lucassen MA; van der Zee R; Voorhorst-Ogink MM; van Kooten PJ; Geuze HJ; van Eden W. Two monoclonal
antibodies generated against human hsp60 show reactivity with synovial membranes of patients with juvenile chronic arthritis. Journal of
Experimental Medicine, 1992, 175(6):1805-10.

Warranty:

No products or "Instructions For Use (IFU)" are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our IFU or website. Our warranty is limited to the actual price paid for the product. ScyTek Laboratories, Inc. is not liable for any property damage, personal injury, time or effort or economic loss caused by our products. Immunohistochemistry is a complex technique involving both histological and immunological detection methods. Tissue processing and handling prior to immunostaining can cause inconsistent results. Variations in fixation and embedding or the inherent nature of the tissue specimen may cause variations in results. Endogenous peroxidase activity or pseudoperoxidase activity in erythrocytes and endogenous biotin may cause non-specific staining depending on detection system used.

Storage: 2° C

ScyTek Laboratories, Inc. 205 South 600 West Logan, UT 84321 U.S.A. CE

EC REP EmergoEurope (31)(0) 70 345-8570 Molsnstraat 15 2513 BH Hague, The Netherlands