

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



Anti-GRP78 Antibody [3G12-1G11]

Mouse Anti-Rat GRP78 Monoclonal IgG1 Kappa Catalog No. SMC-211



Overview

Purification

Product Name
GRP78 Antibody
Description
Mouse Anti-Rat GRP78 Monoclonal IgG1 Kappa
Species Reactivity
Human, Mouse, Rat
Applications
WB, ICC/IF, ELISA
Antibody Dilution
WB (1:1000), ICC/IF (1:100); optimal dilutions for assays should be determined by the user.
Host Species
Mouse
Immunogen Species
Rat
Immunogen
Full-length recombinant rat GRP78
Concentration
1 mg/ml
Conjugates
Alkaline Phosphatase, APC, ATTO 390, ATTO 488, ATTO 565, ATTO 594, ATTO 633, ATTO 655, ATTO 680, ATTO 700, Biotin, FITC, HRP, PE/ATTO 594, PerCP, RPE, Streptavidin, Unconjugated
Properties
Storage Buffer
PBS pH7.4, 50% glycerol, 0.09% sodium azide
Storage Temperature
-20°C
Shipping Temperature
Blue Ice or 4°C

Protein G Purified
Clonality
Monoclonal
Clone Number
3G12-1G11
Isotype
IgG1 Kappa
Specificity
Detects ~78kDa.
Cite This Product
Mouse Anti-Rat GRP78 Monoclonal, Clone 3G12-1G11 (StressMarq Biosciences Inc., Victoria BC CANADA, Catalog # SMC-211)
Certificate Of Analysis
1 μ g/ml of SMC-211 was sufficient for detection of GRP78 in 20 μ g of HEK-293 lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.
Biological Description
Alternative Names 78 kDa glucose regulated protein Antibody, 78 kDa glucose-regulated protein Antibody, AL022860 Antibody, AU019543 Antibody, BIP Antibody, D2Wsu141e Antibody, D2Wsu17e Antibody, Endoplasmic reticulum lumenal Ca(2+)-binding protein grp78 Antibody, Endoplasmic reticulum lumenal Antibody, Ca2+ binding protein grp78 Antibody, FLJ26106 Antibody, Glucose Regu 0 < 2 Antibody
Research Areas
Cancer, Heat Shock, Cell Signaling, Chaperones, Organelle Markers, Trafficking
Cellular Localization
Endoplasmic Reticulum, Endoplasmic reticulum lumen, Melanosome
Accession Number
NP_037215.1
Gene ID
25617
Swiss Prot

Scientific Background

P06761

GRP78 is a ubiquitously expressed, 78-kDa glucose-regulated protein, and is commonly referred to as an immunoglobin chain binding protein (BiP). The BiP proteins are categorized as stress response proteins because they play an important role in the proper folding and assembly of nascent protein and in the scavenging of misfolded proteins in the endoplasmic reticulum lumen. Translation of BiP is directed by an internal ribosomal entry site (IRES) in the 5' nontranslated region of the BiP mRNA. BiP IRES activity increases when cells are heat stressed (1). GRP78 is also critical for maintenance of cell homeostasis and the prevention of apoptosis (2). Luo et al. have provided findings that suggest GRP78 is essential for embryonic cell growth and pluripotent cell survival (3).


In terms of diseases, GRP78 has been shown to be a reliable biomarker of hypoglycemia, to serve a neuroprotective function in

neurons exposed to glutamate and oxidative stress (4), and its protein levels are reduced in the brains of Alzheimers patients (5). Also, the induction of the GRP78 protein that results in severe glucose and oxygen deprivation could possible lead to drug resistance to anti-tumor drugs (6, 7).


References

- 1. Cho, S. et al. (2007). Mol Cell Biol 27(1): 368-83.
- 2. Yang, Y. et al. (1998) J Biol Chem 273: 25552-25555.
- 3. Luo, S. et al (2006) 26 (15): 5688-97.
- 4. Yu, Z. et al. (1999) Exp Neurol. 15: 302-314.
- 5. Koomagi, R. et al. (1999) Anticancer Res. 19: 4333-4336.
- 6. Laquerre, S. et al. (1998) J. Virology 72: 4940-4949.
- 7. Dong, D. et al. (2005) Cancer Res 65(13): 5785-91.

Product Images



Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-GRP78 Monoclonal Antibody, Clone 3G12-1G11 (SMC-211). Tissue: Fibroblast cell line (NIH 3T3). Species: Mouse. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-GRP78 Monoclonal Antibody (SMC-211) at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: DAPI (blue) nuclear stain at 1:5000 for 5 min RT. Localization: Endoplasmic Reticulum, Endoplasmic Reticulum Lumen . Magnification: 60X.



Western Blot analysis of Human, Mouse, Rat HEK-293, NIH3T3, and Rat Brain cell lysates showing detection of GRP78 protein using Mouse Anti-GRP78 Monoclonal Antibody, Clone 3G12-1G11 (SMC-211). Primary Antibody: Mouse Anti-GRP78 Monoclonal Antibody (SMC-211) at 1:1000.

Product Citations (0)

Currently there are no citations for this product.

Reviews

