



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# Anti-ENaC Alpha Antibody [2G4]

Mouse Anti-Rat ENaC alpha Monoclonal IgG2b  
Catalog No. SMC-239



Discovery through partnership | Excellence through quality

## Overview

### Product Name

ENaC alpha Antibody

### Description

Mouse Anti-Rat ENaC alpha Monoclonal IgG2b

### Species Reactivity

Mouse, Rat

### Applications

WB, IHC, IF

### Antibody Dilution

WB (1:1000); IHC (1:150); optimal dilutions for assays should be determined by the user.

### Host Species

Mouse

### Immunogen Species

Rat

### Immunogen

Synthetic peptide from the N-terminal of Rat ENaC alpha (aa. 46-68)

### Concentration

1 mg/ml

### Conjugates

Alkaline Phosphatase, APC, ATTO 390, ATTO 488, ATTO 565, ATTO 594, ATTO 633, ATTO 655, ATTO 680, ATTO 700, Biotin, FITC, HRP, PE/ATTO 594, PerCP, RPE, Streptavidin, Unconjugated

## Properties

### Storage Buffer

PBS pH7.4, 50% glycerol, 0.09% sodium azide

### Storage Temperature

-20°C

### Shipping Temperature

Blue Ice or 4°C

### Purification

Protein G Purified

#### Clonality

Monoclonal

#### Clone Number

2G4

#### Isotype

IgG2b

#### Specificity

Detects ~85kDa.

#### Cite This Product

Mouse Anti-Rat ENaC alpha Monoclonal (StressMarq Biosciences Inc., Victoria BC CANADA, Catalog # SMC-239)

#### Certificate Of Analysis

A 1:1000 dilution of SMC-239 was sufficient for detection of ENaC alpha in 15 µg of Mouse whole kidney lysate by ECL immunoblot analysis using goat anti-mouse IgG:HRP as the secondary antibody.

### Biological Description

#### Alternative Names

SCNN1A Antibody, Epithelial Sodium Channel-? Antibody, Epithelial Sodium Channel alpha Antibody, Alpha ENaC 2 Antibody, Alpha ENaC Antibody, Alpha NaCH Antibody, Alpha-ENaC Antibody, Amiloride sensitive epithelial sodium channel alpha subunit Antibody, Amiloride sensitive sodium channel subunit alpha Antibody, Amiloride-sensitive sodium channel subunit alpha Antibody, ENaCa Antibody, ENaCalpha Antibody, Epithelial Na(+) channel subunit alpha Antibody, Epithelial Na+ channel subunit alpha Antibody, FLJ21883 Antibody, Nonvoltage gated sodium channel 1 subunit alpha Antibody, Nonvoltage-gated sodium channel 1 subunit alpha Antibody, SCNEA Antibody, SCNN 1 Antibody, SCNN1 Antibody, SCNN1A Antibody, SCNNA\_HUMAN Antibody, Sodium channel nonvoltage gated 1 alpha Antibody

#### Research Areas

Epithelial Sodium Channels (ENaC), Ion Channels, Neuroscience, Sodium Channels

#### Cellular Localization

Apical cell membrane

#### Accession Number

NP\_113736

#### Gene ID

25122

#### Swiss Prot

Q6IRJ1

#### Scientific Background

The Epithelial Sodium Channel (ENaC) is a membrane ion channel permeable to Na<sup>+</sup> ions. It is located in the apical plasma membrane of epithelia in the kidneys, lung, colon, and other tissues where it plays a role in trans epithelial Na<sup>+</sup>-ion transport (1). Specifically Na<sup>+</sup> transport via ENaC occurs across many epithelial surfaces, and plays a key role in regulating salt and water absorption (2).


ENaCs are composed of three structurally related subunits that form a tetrameric channel, alpha, beta, and gamma. The

expression of its alpha and beta subunits is enhanced as keratinocytes differentiate (3, 4). The beta and gamma-ENaC subunits are essential for edema fluid to exert its maximal effect on net fluid absorption by distal lung epithelia(5). And it has been concluded that the subunits are differentially expressed in the retina of mice with ocular hypertension, therefore the up-regulation of alpha-ENaC proteins could serve as a protection mechanism against elevated intraocular pressure (6).


## References

1. Kakizoe Y., et al. (2009) J Hypertens. 27(8): 1679-1689.
2. Gu Y. (2008) J Cell Physiol. 216(2):453-457.
3. Bruns J.B. (2003) Am J Physiol Renal Physiol. 285(4): F600-F609.
4. Mauro T., et al. (2002) J Invest Dermatol. 118(4): 589-594.
5. Elias N., et al. (2007) Am J Physiol Lung Cell Mol Physiol. 293(3): L537-45.
6. Dyka F.M., May C.A. and Enz R. (2005) J Neurochem. 94(1): 120-128.


## Product Images



Immunohistochemistry analysis using Mouse Anti-ENaC alpha Monoclonal Antibody, Clone 2G4 (SMC-239). Tissue: Kidney. Species: Rat. Fixation: Paraffin-embedded formalin-fixed. Primary Antibody: Mouse Anti-ENaC alpha Monoclonal Antibody (SMC-239) at 1:100. Secondary Antibody: Goat Anti-Mouse ATTO 488 (green). Localization: Intercalated cells. Aquaporin 2 Antibody staining in red.



Immunohistochemistry analysis using Mouse Anti-ENaC alpha Monoclonal Antibody, Clone 2G4 (SMC-239). Tissue: Kidney (cortex). Species: Mouse. Primary Antibody: Mouse Anti-ENaC alpha Monoclonal Antibody (SMC-239) at 1:150. Localization: Collecting duct principal cells. Magnification: 60X.



Western Blot analysis of Mouse Whole kidney homogenates showing detection of ~85kDa ENaC alpha protein using Mouse Anti-ENaC alpha Monoclonal Antibody, Clone 2G4 (SMC-239). Lane 1: Molecular Weight Ladder (MW). Lane 2: Low-salt diet. Lane 3: Normal-salt diet. Load: 20 µg. Primary Antibody: Mouse Anti-ENaC alpha Monoclonal Antibody (SMC-239) at 1:1000. Predicted/Observed Size: ~85kDa.

## Product Citations (0)

---

Currently there are no citations for this product.

## Reviews

---

There are no reviews yet.