



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

RECOMBINANT  
Anti-Amyloid  
Beta 1-42  
Oligomer  
Antibody



Discovery through Partnership | Excellence through Quality

Rabbit Anti-Human Amyloid Beta 1-42 Oligomer  
Recombinant Monoclonal IgG  
Catalog No. SMC-618

### Product Name

Amyloid Beta 1-42 Oligomer Antibody

### Description

Rabbit Anti-Human Amyloid Beta 1-42 Oligomer Recombinant Monoclonal IgG

### Species Reactivity

Human

### Applications

WB, ELISA

### Antibody Dilution

WB, ELISA (1:1000 - 1:4000); optimal dilutions for assays should be determined by the user.

### Host Species

Rabbit

### Immunogen Species

Human

### Immunogen

Amyloid beta 1-42 oligomers (StressMarq Catalog# SPR-488)

### Concentration

1mg/mL

### Conjugates

APC, ATTO 390, ATTO 488, ATTO 594, Biotin, FITC, HRP, PerCP, RPE, Unconjugated

### Field Of Use

Not for use in humans. Not for use in diagnostics or therapeutics. For in vitro research use only.

## Properties

---

### Storage Buffer

PBS pH 7.4, 50% glycerol, 0.09% sodium azide. \*Storage buffer may change when conjugated

---

### Storage Temperature

-20°C

---

### Shipping Temperature

Blue Ice or 4°C

---

### Purification

Protein A purified

---

### Clonality

Recombinant Monoclonal

---

### Clone Number

1A2

---

### Isotype

IgG

---

### Specificity

Amyloid Beta 1-42 oligomers

---

### Cite This Product

Rabbit Anti-Human Amyloid Beta 1-42 Oligomer Recombinant Monoclonal (StressMarq Biosciences, Victoria BC, Cat# SMC-618)

---

### Certificate Of Analysis

A 1:4000 dilution of SMC-618 was sufficient for detection of 2ng of AB 1-42 oligomers by dot blot analysis with goat-anti-rabbit:HRP as the secondary and by self-sandwich ELISA analysis (capture at 1 ug/mL, detection at 0.2 ug/mL).

---

## Biological Description

---

### Alternative Names

Abeta Oligomers Antibody, APP Antibody

## Research Areas

Alzheimer's Disease, Amyloid, Neurodegeneration, Neuroscience

## Cellular Localization

Membrane (Secreted)

## Gene ID

351 (APP)

## Swiss Prot

P05067


## Scientific Background

Our amyloid beta oligomer antibody has been tested for specificity against AB 1-42 aggregates, with higher specificity to oligomers demonstrated. In the brain, amyloid beta peptide (A $\beta$ ) is generated by protease cleavage of amyloid precursor protein (APP), which aggregates into oligomers, protofibrils, fibrils and ultimately plaques in neurodegenerative diseases. The accumulation of A $\beta$  plaques in the brain is considered a hallmark of Alzheimer's disease (AD), and most of the drugs tested for AD in the past 20 years have targeted amyloid beta accumulation (1). The immunogen used was our Amyloid Beta 1-42 (A $\beta$ 42) Oligomers are generated from Amyloid Beta Peptide 1-42 pre-treated with 1,1,1,3,3,3-Hexafluoro-2-propanol (HFIP) as previously published (2,3).


## References

1. Panza et al. 2019. Nat Rev Neurol. 15:73-88 <https://doi.org/10.1038/s41582-018-0116-6>
2. Stine et al. 2003. JBC. 278(13):11612-22. doi: 10.1074/jbc.M210207200
3. Ahmed et al. 2010. Nature Structural & Molecular Biology. 17(5):561 doi: 10.1038/nsmb.1799


## Product Images



Western blot analysis comparing detection of StressMarq's Amyloid Beta 1-42 constructs (SPR-485 = amyloid beta 1-42 peptide; SPR-488 = amyloid beta 1-42 oligomers; SPR-487 = amyloid beta 1-42 pre-formed fibrils; SPR-492 = amyloid beta pyroglutamate 3-42 pre-formed fibrils). Block: 5% skim milk. Primary Antibodies: Rabbit Anti-Amyloid Beta 1-42 oligomer monoclonal antibody (SMC-618, clone 1A2) & Mouse Anti-Amyloid Beta 1-16 monoclonal antibody (Biolegend 803001, clone 6E10), both at 1:1000 for 1/2 hour at RT. Secondary Antibodies: Goat anti-rabbit IgG:HRP or anti-mouse IgG:HRP at 1:4000 for 1/2 hour at RT. Color Development: Chemiluminescent for HRP (Moss) for 1 min in RT. Exposed 1 second.




Sandwich ELISA analysis comparing detection of Amyloid Beta constructs. The range tested for each construct was 0 to 2000 pg/mL, and the standard curve was prepared using amyloid beta 1-42 oligomers (SPR-488). Within this range, no cross-reactivity with monomeric peptide & APP 751 was observed.



Dot blot analysis comparing detection of StressMarq's amyloid beta 1-42 constructs, alpha synuclein wildtype constructs, and tau constructs. Of note is that SMC-618 does not bind APP 751 or other protein constructs such as alpha synuclein and tau. Block: 5% skim milk. Primary Antibodies: Anti-Amyloid Beta 1-42 oligomer monoclonal antibody (SMC-618, clone 1A2), Anti-Amyloid Beta 1-16 monoclonal antibody (Biolegend 803001, clone 6E10), Anti-Alpha Synuclein polyclonal antibody (SPC-800), and Anti-Tau

polyclonal antibody (SPC-801), all incubated for 1/2 hour at RT. Secondary Antibodies: Goat anti-rabbit IgG:HRP or anti-mouse IgG:HRP at 1:4000 for 1/2 hour at RT. Color Development: Chemiluminescent for HRP (Moss) for 1 min in RT. Exposed 1 second.



Dot blot analysis comparing detection of StressMarq's amyloid beta constructs. Concentrations tested are 20 & 2ng, respectively. Block: 5% skim milk. Primary Antibodies: Rabbit Anti-Amyloid Beta 1-42 oligomer monoclonal antibody (SMC-618, clone 1A2) & Mouse Anti-Amyloid Beta 1-16 monoclonal antibody (Biogen 803001, clone 6E10), both at 1:1000 for 1/2 hour at RT. Secondary Antibodies: Goat anti-rabbit IgG:HRP or anti-mouse IgG:HRP at 1:4000 for 1/2 hour at RT. Color Development: Chemiluminescent for HRP (Moss) for 1 min in RT. Exposed 1 second.

## Product Citations

---

## Reviews

---

There are no reviews yet.