



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Anti-HSP90 Antibody

Rabbit Anti-Human HSP90 Polyclonal  
Catalog No. SPC-104



Discovery through partnership | Excellence through quality

## Overview

---

### Product Name

HSP90 Antibody

### Description

Rabbit Anti-Human HSP90 Polyclonal

### Species Reactivity

Human, Mouse, Rat

### Applications

WB, IHC, ICC/IF, IP, ELISA

### Antibody Dilution

WB (1:500), IHC (1:100), ICC/IF (1:100); optimal dilutions for assays should be determined by the user.

### Host Species

Rabbit

### Immunogen Species

Human

### Immunogen

Full length protein HSP90

### Concentration

1 mg/ml

### Conjugates

Alkaline Phosphatase, APC, ATTO 390, ATTO 488, ATTO 565, ATTO 594, ATTO 633, ATTO 655, ATTO 680, ATTO 700, Biotin, FITC, HRP, PE/ATTO 594, PerCP, RPE, Streptavidin, Unconjugated

## Properties

---

### Storage Buffer

PBS pH 7.4, 50% glycerol, 0.09% sodium azide

### Storage Temperature

-20°C

### Shipping Temperature

Blue Ice or 4°C

### Purification

---

Protein A purified

---

### Clonality

Polyclonal

---

### Specificity

Detects ~90kDa.

---

### Cite This Product

Rabbit Anti-Human HSP90 Polyclonal (StressMarq Biosciences Inc., Victoria BC CANADA, Catalog # SPC-104)

---

### Certificate Of Analysis

A 1:500 dilution of SPC-104 was sufficient for detection of 0.2 mg of purified HSP90 by ECL immunoblot analysis.

---

## Biological Description

---

### Alternative Names

HSP84 Antibody, HSP86 Antibody, HSP90A Antibody, HSP90AA1 Antibody, HSP90AB1 Antibody, HSP90B Antibody, HSPC1 Antibody, HSPC2 Antibody, HSPCAL1 Antibody, HSPCAL4 Antibody, HSP90N Antibody

---

### Research Areas

Cancer, Heat Shock

---

### Cellular Localization

Cytoplasm, Melanosome

---

### Accession Number

NP\_031381.2

---

### Gene ID

3326

---

### Swiss Prot

P08238

---

### Scientific Background

HSP90 is a highly conserved and essential stress protein that is expressed in all eukaryotic cells. From a functional perspective, HSP90 participates in the folding, assembly, maturation, and stabilization of specific proteins as an integral component of a chaperone complex (1-4). Despite its label of being a heat-shock protein, HSP90 is one of the most highly expressed proteins in unstressed cells (1-2% of cytosolic protein). It carries out a number of housekeeping functions – including controlling the activity, turnover, and trafficking of a variety of proteins. Most of the HSP90-regulated proteins that have been discovered to date are involved in cell signaling (5-6). The number of proteins now known to interact with HSP90 is about 100. Target proteins include the kinases v-Src, Wee1, and c-Raf, transcriptional regulators such as p53 and steroid receptors, and the polymerases of the hepatitis B virus and telomerase.5. When bound to ATP, HSP90 interacts with co-chaperones Cdc37, p23, and an assortment of immunophilin-like proteins, forming a complex that stabilizes and protects target proteins from proteasomal degradation. In most cases, HSP90-interacting proteins have been shown to co-precipitate with HSP90 when carrying out immunoadsorption studies, and to exist in cytosolic heterocomplexes with it. In a number of cases, variations in HSP90 expression or HSP90 mutation has been shown to degrade signaling function via the protein or to impair a specific function of the protein (such as steroid binding, kinase activity) in vivo. Ansamycin antibiotics, such as geldanamycin and radicicol, inhibit HSP90 function (7). Looking for more information on HSP90? Visit our new HSP90 Scientific Resource Guide at <http://www.HSP90.ca>.


---

### References

1. Arlander S.J.H., et al. (2003) J Biol Chem 278: 52572-52577.
2. Pearl H., et al. (2001) Adv Protein Chem 59:157-186.



3. Neckers L., et al. (2002) Trends Mol Med 8:S55-S61.
4. Pratt W., Toft D. (2003) Exp Biol Med 228:111-133.
5. Pratt W., Toft D. (1997) Endocr Rev 18: 306360.
6. Pratt W.B. (1998) Proc Soc Exptl Biol Med 217: 420434.
7. Whitesell L., et al. (1994) Proc Natl Acad Sci USA 91: 83248328.

## Product Images



Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-Hsp90 Polyclonal Antibody (SPC-104). Tissue: Heat Shocked HeLa Cells. Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Rabbit Anti-Hsp90 Polyclonal Antibody (SPC-104) at 1:100 for 12 hours at 4°C. Secondary Antibody: FITC Goat Anti-Rabbit (green) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Cytoplasm. Magnification: 100x. (A) DAPI (blue) nuclear stain. (B) Anti-Hsp90 Antibody. (C) Composite. Heat Shocked at 42°C for 1h.

Western blot analysis of Human Cell line lysates showing detection of HSP90 protein using Rabbit Anti-HSP90 Polyclonal Antibody (SPC-104). Load: 2 µg Hsp90beta. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Rabbit Anti-HSP90 Polyclonal Antibody (SPC-104) at 1:10000 for 2 hours at RT. Secondary Antibody: Donkey Anti-Rabbit IgG: HRP for 1 hour at RT.



Immunohistochemistry analysis using Rabbit Anti-HSP90 Polyclonal Antibody (SPC-104). Tissue: backskin. Species: Mouse. Fixation: Bouin's Fixative Solution. Primary Antibody: Rabbit Anti-HSP90 Polyclonal Antibody (SPC-104) at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Rabbit (green) at 1:50 for 1 hour at RT. Localization: Cytoplasm.



Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-Hsp90 Polyclonal Antibody (SPC-104). Tissue: Heat Shocked HeLa Cells. Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Rabbit Anti-Hsp90 Polyclonal Antibody (SPC-104) at 1:100 for 12 hours at 4°C. Secondary Antibody: APC Goat Anti-Rabbit (red) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Cytoplasm. Magnification: 20x. (A) DAPI (blue) nuclear stain. (B) Anti-Hsp90 Antibody. (C) Composite. Heat Shocked at 42°C for 1h.

## Product Citations (3)

---

### Western Blot

---

#### **B Cell Specific Deficiencies in mTOR Limit Humoral Immune Responses.**

Zhang, S. et al. (2013) J Immunol. 191(4):1692-703.

**PubMed ID:** 23858034 **Reactivity:** Mouse **Applications:** Western Blot

#### **Systemic anti-vascular endothelial growth factor therapies induce a painful sensory neuropathy.**

Verheyen, A. et al. (2012) Brain. 135 (9): 2629-2641.

**PubMed ID:** 22734125 **Reactivity:** Mouse **Applications:** Western Blot

#### **Pharmacological inhibition of HSP90 activity negatively modulates myogenic differentiation and cell survival in C2C12 cells.**

Wagatsuma, A. et al. (2011) Mol Cell Biochem. 358 (1-2): 265-280.

**PubMed ID:** 21739150 **Reactivity:** Mouse **Applications:** Western Blot

## Reviews

---

Based on validation through cited publications.

★★★★★

**StressMarq Biosciences**  
June 15, 2016: