



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-ENaC Alpha Antibody

Rabbit Anti-Rat ENaC Alpha Polyclonal
Catalog No. SPC-403



Discovery through partnership | Excellence through quality

Overview

Product Name

ENaC Alpha Antibody

Description

Rabbit Anti-Rat ENaC Alpha Polyclonal

Species Reactivity

Mouse, Rat, African clawed frog (*Xenopus laevis*)

Applications

WB, IHC, ICC/IF

Antibody Dilution

WB (1:1000), ICC/IF (1:400); optimal dilutions for assays should be determined by the user.

Host Species

Rabbit

Immunogen Species

Rat

Immunogen

Produced against a synthetic peptide mapping to the N-terminal of the alpha subunit (amino acids 46-68) of rat Alpha ENaC (antibody designation 3560-2).

Concentration

1 mg/ml

Conjugates

Alkaline Phosphatase, APC, ATTO 390, ATTO 488, ATTO 565, ATTO 594, ATTO 633, ATTO 655, ATTO 680, ATTO 700, Biotin, FITC, HRP, PE/ATTO 594, PerCP, RPE, Streptavidin, Unconjugated

Properties

Storage Buffer

PBS, 50% glycerol, 0.09% sodium azide

Storage Temperature

-20°C

Shipping Temperature

Blue Ice or 4°C

Purification

Protein A purified

Clonality

Polyclonal

Specificity

Detects ~85kDa.

Cite This Product

Rabbit Anti-Rat ENaC Polyclonal (StressMarq Biosciences Inc., Victoria BC CANADA, Catalog # SPC-403)

Certificate Of Analysis

1 µg/ml of SPC-403 was sufficient for detection of alpha-ENaC in 35 µg of rat kidney tissue lysate by colorimetric immunoblot analysis using Goat anti-rabbit IgG:HRP as the secondary antibody.

Biological Description

Alternative Names

SCNN1A Antibody, Epithelial Sodium Channel-? Antibody, Epithelial Sodium Channel alpha Antibody, Alpha ENaC 2 Antibody, Alpha ENaC Antibody, Alpha NaCH Antibody, Alpha-ENaC Antibody, Amiloride sensitive epithelial sodium channel alpha subunit Antibody, Amiloride sensitive sodium channel subunit alpha Antibody, Amiloride-sensitive sodium channel subunit alpha Antibody, ENaCa Antibody, ENaCalpha Antibody, Epithelial Na(+) channel subunit alpha Antibody, Epithelial Na+ channel subunit alpha Antibody, FLJ21883 Antibody, Nonvoltage gated sodium channel 1 subunit alpha Antibody, Nonvoltage-gated sodium channel 1 subunit alpha Antibody, SCNEA Antibody, SCNN 1 Antibody, SCNN1 Antibody, SCNN1A Antibody, SCNNA_HUMAN Antibody, Sodium channel nonvoltage gated 1 alpha Antibody

Research Areas

Epithelial Sodium Channels (ENaC), Ion Channels, Neuroscience, Sodium Channels

Cellular Localization

Apical cell membrane

Accession Number

NP_113736

Gene ID

25122

Swiss Prot

Q6IRJ1

Scientific Background


The Epithelial Sodium Channel (ENaC) is a membrane ion channel permeable to Na⁺ ions. It is located in the apical plasma membrane of epithelia in the kidneys, lung, colon, and other tissues where it plays a role in trans epithelial Na⁺-ion transport (1). Specifically Na⁺ transport via ENaC occurs across many epithelial surfaces, and plays a key role in regulating salt and water absorption (2).

ENaCs are composed of three structurally related subunits that form a tetrameric channel, α , β , and γ . The expression of its alpha and beta subunits is enhanced as keratinocytes differentiate (3, 4). The beta and gamma-ENaC subunits are essential for edema fluid to exert its maximal effect on net fluid absorption by distal lung epithelia(5). And it has been concluded that the subunits are differentially expressed in the retina of mice with ocular hypertension, therefore the up-regulation of alpha-ENaC proteins could serve as a protection mechanism against elevated intraocular pressure (6).


References

1. Kakizoe Y., et al. (2009) *J Hypertens.* 27(8): 1679-1689.
2. Gu Y. (2008) *J Cell Physiol.* 216(2):453-457.
3. Bruns J.B. (2003) *Am J Physiol Renal Physiol.* 285(4): F600-F609.
4. Mauro T., et al. (2002) *J Invest Dermatol.* 118(4): 589-594.
5. Elias N., et al. (2007) *Am J Physiol Lung Cell Mol Physiol.* 293(3): L537-45.
6. Dyka F.M., May C.A. and Enz R. (2005) *J Neurochem.* 94(1): 120-128.

Product Images



Western blot analysis of Rat kidney tissue lysates showing detection of ENaC protein using Rabbit Anti-ENaC Polyclonal Antibody (SPC-403). Primary Antibody: Rabbit Anti-ENaC Polyclonal Antibody (SPC-403) at 1:1000.



Western blot analysis of Mouse kidney tissue lysates showing detection of ENaC protein using Rabbit Anti-ENaC Polyclonal Antibody (SPC-403). Primary Antibody: Rabbit Anti-ENaC Polyclonal Antibody (SPC-403) at 1:1000.

Product Citations (26)

Western Blot

Renal sodium transport in renin-deficient Dahl salt-sensitive rats.

Pavlov, T.S., Levchenko, V., Ilatovskaya, D.V., Moreno, C. and Staruschenko, A. (2016) *J Renin Angiotensin Aldosterone Syst.* 17(3).

PubMed ID: 27443990 **Reactivity:** Rat **Applications:** Western Blot

(Pro)Renin Receptor Regulates Potassium Homeostasis through a Local Mechanism.

Xu, C. et al. (2016) *Am J Physiol Renal Physiol.* [Epub ahead of print]

PubMed ID: 27440776 **Reactivity:** Rat **Applications:** Western Blot

AMP-activated protein kinase attenuates high salt-induced activation of epithelial sodium channels (ENaC) in human umbilical vein endothelial cells.

Zheng, W. et al. (2016) *Oxid Med Cell Longev*. [Epub ahead of print]

PubMed ID: **Reactivity:** Human **Applications:** Western Blot

Renal tubular epithelial cell prorenin receptor regulates blood pressure and sodium transport.

Ramkumar, N. et al. (2016) *Am J Physiol Renal Physiol*. [Epub ahead of print]

PubMed ID: 27053687 **Reactivity:** Mouse **Applications:** Western Blot

The Role of Epithelial Sodium Channel ENaC and the Apical Cl⁻/HCO₃⁻ Exchanger Pendrin in Compensatory Salt Reabsorption in the Setting of Na-Cl Cotransporter (NCC) Inactivation.

Patel-Chamberlin, M. et al. (2016) *PLoS One*. 11(3):e0150918.

PubMed ID: 26963391 **Reactivity:** Mouse **Applications:** Western Blot

Possible role for nephron-derived angiotensinogen in angiotensin-II dependent hypertension.

Ramkumar, N. et al. (2016) *Physiol Rep*. 4(1). pii: e12675.

PubMed ID: 26755736 **Reactivity:** Mouse **Applications:** Western Blot

The Thiazide-Sensitive Co-Transporter Promotes the Development of Sodium Retention in Mice with Diet-Induced Obesity.

Davies, M. R. P. et al. (2015) *Kidney Blood Press Res*. 40(5):509-519.

PubMed ID: 26418861 **Reactivity:** Mouse **Applications:** Western Blot

Inhibition of Mitochondrial Complex-1 Prevents the Downregulation of NKCC2 and ENaC? in Obstructive Kidney Disease.

Zhang, Y. et al. (2015) *Sci Rep*. 5:12480.

PubMed ID: 26207612 **Reactivity:** Mouse **Applications:** Western Blot

Increased Epithelial Sodium Channel Activity Contributes to Hypertension Caused by Na⁺-HCO₃⁻ Cotransporter Electrogenic 2 Deficiency.

Wen, D. et al. (2015) *Hypertension*. 66(1):68-74.

PubMed ID: 25941340 **Reactivity:** Mouse **Applications:** Western Blot

Compositions And Methods Of Use For (PRO)Renin Receptor.

Feng, Y. and Yang, T. (2015) United States Patent Application 20150025013.

PubMed ID: **Reactivity:** Rat **Applications:** Western Blot

Role of Rho GDP-dissociation inhibitor alpha in control of ENaC-mediated sodium reabsorption.

Pavlov, T. S., Levchenko, V. and Staruschenko, A. (2014) *J Biol Chem*. 289(41):28651-9.

PubMed ID: 25164814 **Reactivity:** Mouse **Applications:** Western Blot

Collecting duct specific knock-out of renin attenuates angiotensin-II induced hypertension.

Ramkumar, N. et al. (2014) *Am J Physiol Renal Physiol*. 307(8):F931-8.

PubMed ID: 25122048 **Reactivity:** Mouse **Applications:** Western Blot

Novel Mechanisms of Sodium Retention in Obesity: Phosphorylation of NKCC2 and Regulation of SPAK/OSR1 by AMPK.

Davies, M. R. P. et al. (2014) *Am J Physiol*. 307(1):F96-F106.

PubMed ID: 24808538 **Reactivity:** Mouse **Applications:** Western Blot

Chronic Angiotensin II Infusion Drives Extensive Aldosterone-Independent Epithelial Na⁺ Channel Activation.

Mamenko, M. et al. (2013) *Hypertension*. 62(6):1111-22.

PubMed ID: 24060890 **Reactivity:** Mouse **Applications:** Western Blot

Adenylyl Cyclase VI Mediates Vasopressin-Stimulated ENaC Activity.

Roos, K.P. et al. (2012) JASN. 24 (2): 218-227.

PubMed ID: 23264685 **Reactivity:** Mouse **Applications:** Western Blot

Increased expression of renal TRPM6 compensates for Mg²⁺ wasting during furosemide treatment.

van Angelen, A.A., Kemp, A.W., Hoenderop, J.G. and Bindels, B.J. (2012) Clin Kidney J. 5 (6): 535-544.

PubMed ID: 26069797 **Reactivity:** Mouse **Applications:** Western Blot

Regional differences in rat conjunctival ion transport activities.

Yu, D. et al. (2012) Am J Physiol Cell Physiol. 303 (7): 767-780.

PubMed ID: 22814399 **Reactivity:** Xenopus laevis **Applications:** Western Blot

The Epithelial Sodium Channel (ENaC) Establishes a Trafficking Vesicle Pool Responsible for Its Regulation.

Edinger, R.S. et al. (2012) PLoS ONE. 7(9): e46593.

PubMed ID: 23029554 **Reactivity:** Mouse **Applications:** Western Blot

Aldosterone does not require angiotensin II to activate NCC through a WNK4/SPAK-dependent pathway.

van der Lubbe, N. et al. (2012) Euro J Physiol. 463 (6): 853-863.

PubMed ID: 22549242 **Reactivity:** Rat **Applications:** Western Blot

Immunohistochemistry

Inhibition of Mitochondrial Complex-1 Prevents the Downregulation of NKCC2 and ENaC^v in Obstructive Kidney Disease.

Zhang, Y. et al. (2015) Sci Rep. 5:12480.

PubMed ID: 26207612 **Reactivity:** Mouse **Applications:** Immunohistochemistry

Epoxyeicosatrienoic Acid (EET) Analog Lowers Blood Pressure Through Vasodilation And Sodium Channel Inhibition.

Khan, M. A., et al. (2014) Clin Sci (Lond). 127(7):463-74.

PubMed ID: 24707975 **Reactivity:** Rat **Applications:** Immunohistochemistry

Chronic Angiotensin II Infusion Drives Extensive Aldosterone-Independent Epithelial Na⁺ Channel Activation.

Mamenko, M. et al. (2013) Hypertension. 62(6):1111-22.

PubMed ID: 24060890 **Reactivity:** Mouse **Applications:** Immunohistochemistry

ENaC^v-expressing astrocytes in the circumventricular organs, white matter, and ventral medullary surface: sites for Na⁺ regulation by glial cells.

Miller, R.L. and Loewy, A.D. (2013) J Chem Neuroanat. 53:72-80.

PubMed ID: 24145067 **Reactivity:** Rat **Applications:** Immunohistochemistry

Regulation of ENaC in mice lacking renal insulin receptors in the collecting duct.

Pavlov, T.S. et al. (2013) FASEB J. 27(7):2723-32.

PubMed ID: 23558339 **Reactivity:** Mouse **Applications:** Immunohistochemistry

Other Citations

mTORC1 maintains renal tubular homeostasis and is essential in response to ischemic stress.

Grahammer, F. et al. (2014) Proc Natl Acad Sci U S A. 111(27):E2817-26.

PubMed ID: 24958889 **Reactivity:** Mouse **Applications:** Gel Shift

mPGES-1-Derived PGE₂ Mediates Dehydration Natriuresis.

Jia, Z. et al. (2013) Am J Physiol Renal Physiol. 304(2):F214-21.

Reviews

Based on validation through cited publications.



StressMarq Biosciences

June 15, 2016: