



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Amyloid Beta Protein



Discovery through Partnership | Excellence through Quality

Human Synthetic Amyloid Beta 1-42 Oligomers
Catalog No. SPR-488

Product Name

Amyloid Beta Protein

Description

Human Synthetic Amyloid Beta 1-42 Oligomers

Applications

WB, In vivo Assay, In vitro Assay

Concentration

Lot/batch specific. See included datasheet.

Conjugates

No tag

Nature

Synthetic (TFA preparation)

Species

Human

Expression System

N/A

Amino Acid Sequence

DAEFRHDSGYEVHHQKLVFFAEDVGSNKGAIIGLMVGGVVIA

Purity

>95%

Protein Length

42 amino acids

Field Of Use

Not for use in humans. Not for use in diagnostics or therapeutics. For in vitro research use only.

Properties

Storage Buffer

Phosphate buffer (PB) pH 7.4 and 10 mM NaCl

Storage Temperature

-80°C

Shipping Temperature

Dry Ice. Shipping note: Product will be shipped separately from other products purchased in the same order.

Purification

N/A

Cite This Product

Human Synthetic Amyloid Beta Oligomers (StressMarq Biosciences Inc., Victoria BC CANADA, Catalog # SPR-488)

Certificate Of Analysis

Certified >95% pure using mass spec and HPLC.

Biological Description

Alternative Names

Abeta Oligomers, Abeta peptide, Amyloid beta peptide oligomers, Beta amyloid peptide oligomers, amyloid beta precursor protein peptide oligomers, APP

Research Areas

Alzheimer's Disease, Amyloid, Neurodegeneration, Neuroscience

Cellular Localization

Cell membrane, Intracellular Vesicles

Gene ID

351

Swiss Prot

P05067


Scientific Background

Our Amyloid Beta 1-42 (A β 42) Oligomers are generated from Amyloid Beta Peptide 1-42 pre-treated with 1,1,1,3,3,3-Hexafluoro-2-propanol (HFIP) as previously published (1,2). Our A β 42 oligomers present as globular oligomers when observed under TEM and AFM, and have a unique dimer/trimer and oligomer signal on a Western Blot with an anti-amyloid beta antibody. Our A β 42 oligomers were also demonstrated to be toxic to primary rat cortical neurons in a dose-dependent manner. In the brain, amyloid beta peptide (A β) is generated by protease cleavage of amyloid precursor protein (APP), which aggregates into oligomers, protofibrils, fibrils and ultimately plaques in neurodegenerative diseases. The accumulation of A β plaques in the brain is considered a hallmark of Alzheimer's disease (AD), and most of the drugs tested for AD in the past 20 years have targeted amyloid beta accumulation (3). Soluble A β oligomers isolated from the brains of AD patients or those generated in vitro potently impaired synapse structure and function (4). A β oligomers generated in vitro were toxic to PC12 cells (5) and SH-SY5Y cells (6). A β was demonstrated to interact with tauopathies to affect neurodegeneration in AD patients (7) and accumulations of A β were shown to be associated with lower survival rates in Parkinson's disease patients with dementia (8).


References

1. Stine et al. 2003. JBC. 278(13):11612-22. doi: 10.1074/jbc.M210207200
2. Ahmed et al. 2010. Nature Structural & Molecular Biology. 17(5):561-7. doi: 10.1038/nsmb.1799
3. Panza et al. 2019. Nat Rev Neurol. 15:73-88 <https://doi.org/10.1038/s41582-018-0116-6>
4. Shankar et al. 2008. Nat Med. 14(8):837-842. doi: 10.1038/nm1782
5. Chromy et al. 2003. Biochemistry. 42:12749-12760. doi: 10.1021/bi030029q
6. Kaye et al. 2003. Science. 300(5618): 486-489. doi: 10.1126/science.1079469
7. Want et al. 2016. JAMA Neurol. 73(9):1070-7. doi: 10.1001/jamaneurol.2016.2078
8. Kotzbauer et al. 2012. Arch Neurol. 69(10): 1326-1331. doi: 10.1001/archneurol.2012.1608


Product Images




TEM of amyloid beta 1-42 monomers (SPR-485, left), oligomers (SPR-488, middle) and fibrils (SPR-487, right). Negative stain transmission electron microscopy images acquired at 80 Kv on carbon coated 400 mesh copper grids using phosphotungstic acid and uranyl acetate stain. Scale bar = 100 nm.



AFM of amyloid beta 1-42 monomers (SPR-485, left), oligomers (SPR-488, middle) and fibrils (SPR-487, right). Atomic force microscopy analysis of 1.0 mg/mL samples diluted to 0.1 mg/mL in dH₂O, mounted on freshly cleaved mica, washed, dried and analyzed with tapping mode. Representative images are 2.5 x 2.5 μ m x-y with a z-range of 10 nm.



Western blot of amyloid beta 1-42 monomers (SPR-485, left), oligomers (SPR-488, middle) and fibrils (SPR-487, right) using anti-amyloid beta 6E10 antibody. Amyloid beta constructs at 160 pmol were run on 4-12% Bis-Tris SDS-PAGE, transferred to nitrocellulose in the presence of 0.02% v/v Tween-20, and blotted with 1:1000 mouse 6E10 primary antibody (Biolegend). Oligomers observed under TEM/AFM show distinct dimer/trimer bands as well as a signal from ~37-75 kDa (middle). Fibrils observed under TEM/AFM show a signal greater than 100 kDa and a distinct signal in the stacking gel (right).



Amyloid beta 1-42 oligomers (SPR-488) and fibrils (SPR-487) show a dose-dependent toxicity to primary rat cortical neurons, but not monomers (SPR-485). Survival of rat primary cortical neurons 14 days after treatment with different concentrations of (A) monomers, (B) oligomers or (C) fibrils quantified by MAP2 positive neurons and expressed as a percentage of control. Fibrils and respective vehicle controls were initially sonicated in a Bioruptor. Test conditions were run in the same plate as untreated control and vehicle controls, which consisted of buffer without amyloid beta 1-42 protein. Data expressed as mean \pm s.e.m. (n=6). A global analysis of the data was performed using a one-way ANOVA followed by Dunnett's test; ** $p < 0.01$ stats vs control; ## $p < 0.01$, #### $p < 0.0001$ stats vs vehicle control. § represents untreated control condition.

Product Citations (0)

Currently there are no citations for this product.

Reviews

There are no reviews yet.