



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Bosutinib

Chemical Properties

CAS No. :	380843-75-4
Formula:	C ₂₆ H ₂₉ Cl ₂ N ₅ O ₃
Molecular Weight:	530.45
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Bosutinib (SKI-606) is a synthetic quinolone derivative and dual kinase inhibitor that targets both Abl (IC ₅₀ : 1 nM) and Src (IC ₅₀ : 1.2 nM) kinases.
Targets(IC ₅₀)	Bcr-Abl,Src,Autophagy
In vitro	Bosutinib has the antiproliferative activity against three different Bcr-Abl-positive leukemia cell lines (KU812, K562, and MEG-01). Bosutinib inhibited the proliferation of all three cell lines, with IC ₅₀ s ranging from 5 nM in the KU812 line to 20 nM for the K562 and MEG-01 cell lines. Inhibition of proliferation by Bosutinib is associated with cell cycle arrest and cell death. Treatment with Bosutinib at 100 nM for 24 h (KU812) or 48 h (K562) resulted in a reduction of S and G ₂ -M phase cells and an increase of cells with a DNA content of less than 2N. Treatment with Bosutinib at 100 nm also led to PARP degradation after 48 h. The potent antiproliferative activity of Bosutinib against CML lines was not a general property for leukemia cell lines. Molt-4, HL-60, Ramos, and other leukemia cell lines were unaffected by Bosutinib at concentrations less than 1 μM [2].
In vivo	Bosutinib (30/25 mg/kg, b.i.d) reduces tumor growth in unstaged and staged Src-transformed fibroblast mouse xenograft models. Bosutinib (100 mg/kg) also induces complete tumor regression in a K562 mouse xenograft model when administered once per day for five days [2].
Kinase Assay	The Src kinase activity is measured in an ELISA format. Src (3 units/reaction), reaction buffer (50 mM Tris-HCl pH 7.5, 10 mM MgCl ₂ , 0.1 mM EGTA, 0.5 mM Na ₃ VO ₄) and cdc2 substrate peptide are added to various concentration of Bosutinib and incubated at 30 °C for 10 minutes. The reaction is started by the addition of ATP to a final concentration of 100 μM, incubated at 30 °C for 1 hour and stopped by addition of EDTA. Instructions from the manufacturer are followed for subsequent steps. The Abl kinase assay is performed in a DELFIA solid phase europium-based detection assay format. Biotinylated peptide (2 μM) is bound to streptavidin-coated microtitration plates for 1.5 hours in 1 mg/mL ovalbumin in PBS. The plates are washed for 1 hour with PBS/0.1% Tween 80, followed by a PBS wash. The kinase reaction is incubated for 1 hour at 30°C. Abl kinase (10 units) is mixed with 50 mM Tris-HCl (pH 7.5), 10 mM MgCl ₂ , 80 μM EGTA, 100 μM ATP, 0.5 mM Na ₃ VO ₄ , 1% DMSO, 1 mM HEPES (pH 7.0), 200 μg/mL ovalbumin and various concentration of Bosutinib. The reaction is stopped with EDTA at a final concentration of 50 mM. The reaction is monitored with Eu-labeled phosphotyrosine antibody and DELFIA enhancement solution [2].

A DRUG SCREENING EXPERT

Cell Research	Cells are exposed to various concentrations of Bosutinib for 72 hours. Anchorage-independent proliferation of Abl-MLV-transformed fibroblasts is measured in 96-well ultra-low binding plates treated with Sigmacote to block residual cell attachment. Cell proliferation is measured with MTS or Cell-Glo. For the determination of cell cycle or cell death, cells are prepared for FACS analysis in the CycleTest Plus DNA reagent kit and analyzed on a fluorescence-activated cell sorter flow cytometer [2].
Animal Research	K562 cells were suspended to 50 million cells/ml in Matrigel (1 volume of cells with 1 volume of cold Matrigel). Nude female mice 6-7 weeks of age were given injections of 0.2 ml of this suspension. Tumors were staged for 10 days, at which time they entered the growth phase. At this time, the compound was administered by oral gavage in a 0.2-ml suspension with 0.5% methocel/0.4% Tween 80 [2].

Solubility Information

Solubility	DMSO: 65 mg/mL (122.54 mM), Ethanol: 13.3 mg/mL (25 mM), (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.8852 mL	9.426 mL	18.8519 mL
5 mM	0.377 mL	1.8852 mL	3.7704 mL
10 mM	0.1885 mL	0.9426 mL	1.8852 mL
50 mM	0.0377 mL	0.1885 mL	0.377 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Cheng S, Jin P, Li H, et al. Evaluation of CML TKI Induced Cardiovascular Toxicity and Development of Potential Rescue Strategies in a Zebrafish Model. *Frontiers in Pharmacology*. 2021: 2866.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481