



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)


[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Sumatriptan succinate

## Chemical Properties

CAS No. :	103628-48-4
Formula:	C <sub>18</sub> H <sub>27</sub> N <sub>3</sub> O <sub>6</sub> S
Molecular Weight:	413.49
Appearance:	no data available
Storage:	Powder: -20°C for 3 years   In solvent: -80°C for 1 year



## Biological Description

Description	Sumatriptan succinate (GR 43175), a serotonin <sub>1</sub> (5-HT <sub>1</sub> ) receptor agonist, is used in the acute treatment of a migraine headache.
Targets(IC <sub>50</sub> )	5-HT Receptor
In vitro	Sumatriptan is rapidly cleared through metabolism and renal elimination, with a half-life of 1-2 hours. It rarely produces adverse effects when administered acutely, except for dogs exhibiting low tolerance to high doses. Clinically, subcutaneous injection of Sumatriptan at 100 mg/kg notably decreases injury-induced and contralateral mechanical allodynia in a rat model of trigeminal neuropathic pain (peak effects at 6.3g and 4.4g, respectively). In cats, Sumatriptan reduces the number of Fos-positive cells in the caudal part of the spinal trigeminal nucleus (I, IIO, and C2) following mechanical stimulation (to 6, 13, and 9 cells, respectively). The compound's biological activity varies across species, with 37%, 23%, and 58% in rats, rabbits, and dogs, respectively. Sumatriptan selectively constricts cranial blood vessels, which are dilated and inflamed during migraines, mediated by the 5-HT <sub>1</sub> receptor subtype in animal cranial vasculature.
In vivo	Sumatriptan significantly reduces plasma protein-induced extravasation triggered by electrical stimulation of the trigeminal nerve. It mitigates morphological changes in the capillaries within the dura mater and small veins in the presence of hypertrophied mast cells stimulated at the trigeminal ganglion. Sumatriptan has a slightly lower affinity for the serotonin receptor 1A binding site (K <sub>i</sub> =100 nM), while exhibiting the highest affinity for 5HT (K <sub>i</sub> =17 nM) and 5HT <sub>1B</sub> receptor binding sites (K <sub>i</sub> =27 nM).

## Solubility Information

Solubility	DMSO: 55 mg/mL (133.01 mM), H <sub>2</sub> O: 41.4 mg/mL (100 mM), ( < 1 mg/ml refers to the product slightly soluble or insoluble)
------------	---

### Preparing Stock Solutions

---

	1mg	5mg	10mg
1 mM	2.4184 mL	12.0922 mL	24.1844 mL
5 mM	0.4837 mL	2.4184 mL	4.8369 mL
10 mM	0.2418 mL	1.2092 mL	2.4184 mL
50 mM	0.0484 mL	0.2418 mL	0.4837 mL

---

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

### Reference

Peroutka SJ, et al. Eur J Pharmacol, 1989, 163(1), 133-136.

Buzzi MG, et al. Cephalalgia, 1991, 11(4), 165-168.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481