



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Rosiglitazone

Chemical Properties

CAS No. :	122320-73-4
Formula:	C ₁₈ H ₁₉ N ₃ O ₃ S
Molecular Weight:	357.43
Appearance:	no data available
Storage:	store at low temperature Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Rosiglitazone (BRL49653) is a PPAR γ agonist, TRPC5 activator, and TRPM3 inhibitor with oral activity. Rosiglitazone is also a hypoglycemic agent and a thiazolidinedione insulin sensitizer.
Targets(IC50)	Ferroptosis,TRP/TRPV Channel,PPAR,Autophagy
In vitro	Rosiglitazone reduces bone formation rate and increases adipose content within the bone marrow. It decreases the expression of osteoblast-specific genes Runx2/Cbfa1, DLX5, and α 1(I) collagen, while expression of the adipocyte-specific fatty acid binding protein AP2 is increased. This drug leads to significant bone loss, evidenced by reductions in bone mass, trabecular width, and number, along with an increase in trabecular separation. Furthermore, in ob/ob mice, rosiglitazone enhances transcription of genes encoding mitochondrial proteins in white adipocytes, which is accompanied by changes in mitochondrial number and structure.
In vivo	In certain cell lines, Rosiglitazone reduces cholesterol synthesis independent of peroxisome proliferator-activated receptor γ (PPAR γ). The compound significantly enhances the phosphorylation of threonine 172 in the α subunit of AMP-dependent protein kinase, increasing the AMP: ATP ratio. Additionally, Rosiglitazone boosts secretion of adiponectin by up to 2.3-fold from omental cells, while secretion from subcutaneous fat cells remains unaffected. In 3T3-L1 adipocytes, Rosiglitazone alters mitochondrial morphological characteristics and protein profile. It activates complexes containing α 1- and α 2-AMPK, leading to a marked increase in phosphorylation of acetyl-CoA carboxylase. Rosiglitazone also acts as a dominant inhibitor of osteoblastogenesis from mouse marrow in vitro through the activation of PPAR-gamma2.
Cell Research	Rosiglitazone is dissolved in DMSO and stored, and then diluted with appropriate medium before use[2]. Human neuroblastoma SH-SY5Y cells are maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum, 100 μ g/mL Streptomycin and 100 U/mL Penicillin G. SH-SY5Y cells are transfected with the longest isoform of human tau (2N4R) tagged with GFP using lipofectamine. 24 hr after transfection, cells are treated with Rosiglitazone (10 μ M, 50 μ M) for 24 hr[2].

Solubility Information

Solubility	1eq. HCl: 35.7 mg/mL (100 mM), DMSO: 45 mg/mL (125.9 mM), (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.7978 mL	13.9888 mL	27.9775 mL
5 mM	0.5596 mL	2.7978 mL	5.5955 mL
10 mM	0.2798 mL	1.3989 mL	2.7978 mL
50 mM	0.056 mL	0.2798 mL	0.5596 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Zeng X, Zhu S, Lu W, et al. Target identification among known drugs by deep learning from heterogeneous networks. *Chemical Science*. 2020, 11(7): 1775-1797.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 36 Washington Street, Wellesley Hills, MA 02481