



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Streptozocin

Chemical Properties

CAS No. :	18883-66-4
Formula:	C ₈ H ₁₅ N ₃ O ₇
Molecular Weight:	265.22
Appearance:	no data available
Storage:	keep away from direct sunlight, keep away from moisture, store under nitrogen, store at low temperature Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Streptozocin (NSC-85998) is an antibiotic that induces DNA methylation. Streptozocin is toxic to insulin-producing pancreatic islet B-cells and is commonly used in the construction of animal models of type 1 diabetes. The product is unstable in solution and is recommended to be dispensed now.
Targets(IC50)	DNA Alkylator/Crosslinker, DNA Alkylation, DNA/RNA Synthesis, Antibacterial, Antibiotic, Autophagy
In vitro	<p>METHODS: Rat β-insulinoma cells Rin-5F were treated with Streptozocin (1-10 mM) for 2-48 h. Cell viability was measured using MTT assay.</p> <p>RESULTS: Maximum inhibition (60-70%) was observed in cells treated with 10 mM Streptozocin for 24 h and 48 h. The cells were treated with Streptozocin (1-10 mM) for 2-48 h, and cell viability was measured by MTT assay. [1]</p> <p>METHODS: Neural stem cells were treated with Streptozocin (2.5 mM) for 2 days and gene expression was detected using RT-qPCR.</p> <p>RESULTS: Streptozocin significantly reduced the relative expression level of GLUT3 mRNA by 46.4%, and did not affect the relative expression level of IR and GLUT1. [2]</p>
In vivo	<p>METHODS: Streptozocin (40 mg/kg in the 50 mM sodium citrate buffer (pH 4.5), ready to use) was administered intraperitoneally to C57BL/6 or CD-1 male mice once daily for five days. Normal food and 10% sucrose water were provided during the administration period, and 10% sucrose water was replaced with normal water on the sixth day.</p> <p>RESULTS: Repeated low doses of Streptozocin induced type 1 diabetes in mice. [3]</p> <p>METHODS: Streptozocin (200 mg/kg in the 50 mM sodium citrate buffer (pH 4.5), ready to use) was administered to C57BL/6 or CD-1 male mice by a single intraperitoneal injection. Normal food and 10% sucrose water were provided after administration, and 10% sucrose water was replaced with normal water on the third day.</p> <p>RESULTS: A single high dose of Streptozocin induced type 1 diabetes in mice. [3]</p>
Cell Research	Streptozocin (STZ) stock solutions of 50 mg/mL in distilled Water (dWater) are freshly prepared for each experiment[2]. Human and murine cell lines are cultured in triplicate in 96-well plates at a density of 2×10^4 cells/well in the absence (untreated control) or presence of various concentrations of ALX (20-3000 μ g/mL) or STZ (1-3000 μ g/mL) for 48 h at 37°C under a humidified atmosphere containing 5% CO ₂ . Cells incubated in complete media including dWater in a final concentration of 0.1% served as control for solvent toxicity and cells incubated in complete medium are used as a control for the

experiments. The effects of the tested drugs on tumor cell growth or viability are determined employing the MTT assay in accordance with the manufacturer's instructions. The IC50 values (drug concentration that induces 50% inhibition of the cell growth) are calculated using the GraphPad Prism 4 program[2].

Solubility Information

Solubility	H2O: 113.3 mg/mL (427.19 mM), Sonication and heating are recommended. DMSO: 16.67 mg/mL (62.84 mM), Sonication is recommended. (The compound is unstable in solution, please use soon.) (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	---

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.7705 mL	18.8523 mL	37.7045 mL
5 mM	0.7541 mL	3.7705 mL	7.5409 mL
10 mM	0.377 mL	1.8852 mL	3.7705 mL
50 mM	0.0754 mL	0.377 mL	0.7541 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Li Y, Zhou X, Cheng C, et al. Gut AstA mediates sleep deprivation-induced energy wasting in *Drosophila*. *Cell Discovery*. 2023, 9(1): 49.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 36 Washington Street, Wellesley Hills, MA 02481