



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Doxycycline

Chemical Properties

CAS No. :	564-25-0
Formula:	C ₂₂ H ₂₄ N ₂ O ₈
Molecular Weight:	444.43
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Doxycycline (Doxiciclina) belongs to the tetracycline class of antibiotics and is a broad-spectrum metalloproteinase (MMP) inhibitor with oral activity. Doxycycline has antibacterial activity and antitumor activity.
Targets(IC50)	MMP, ribosome, Antibacterial, Antibiotic, Parasite
In vitro	<p>METHODS: Twenty-two tumor cells were treated with Doxycycline for 48 h. Cell viability was measured by MTT.</p> <p>RESULTS: Lung cancer cells were more sensitive to Doxycycline than most other cell lines. NCI-H446 and A549 cells were sensitive to Doxycycline with IC50s of 1.70 μM and 1.06 μM, respectively. [1]</p> <p>METHODS: Human pancreatic cancer cells PANC-1 were treated with Doxycycline (10-40 μg/mL) for 8-24 h. The cell cycle was detected by Flow Cytometry.</p> <p>RESULTS: Doxycycline caused the cells to arrest in the G1-S phase of the cell cycle. [2]</p>
In vivo	<p>METHODS: To detect anti-tumor activity in vivo, Doxycycline (50 mg/kg) was administered intraperitoneally once daily for fifteen days to BALB/c mice harboring human mammary carcinoma tumor 4T1.</p> <p>RESULTS: Doxycycline delayed tumor growth and reduced the number of tumor-associated macrophages and blood vessels. [3]</p> <p>METHODS: To assay antimicrobial activity in vivo, Doxycycline (40 mg/kg) was administered intraperitoneally to BALB/c mice exposed to <i>B. pseudomallei</i> 1026b twice daily for two weeks.</p> <p>RESULTS: Doxycycline-treated mice survived 100% of the time, and no <i>B. pseudomallei</i> DNA was amplified from the lungs or spleens of the majority of the surviving mice. [4]</p>

Solubility Information

Solubility	DMSO: 50 mg/mL (112.5 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.2501 mL	11.2504 mL	22.5007 mL
5 mM	0.450 mL	2.2501 mL	4.5001 mL
10 mM	0.225 mL	1.125 mL	2.2501 mL
50 mM	0.045 mL	0.225 mL	0.450 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Tan Y, Chen X, Xu Z, et al. The mechanics of tumor cells dictate malignancy via cytoskeleton-mediated APC/Wnt/ β -catenin signaling. *Research*. 2023
Qin Y, et al. Doxycycline reverses epithelial-to-mesenchymal transition and

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 36 Washington Street, Wellesley Hills, MA 02481