



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Kaempferol

Chemical Properties

CAS No. :	520-18-3
Formula:	C ₁₅ H ₁₀ O ₆
Molecular Weight:	286.24
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Kaempferol (Robigenin) is a natural flavonoid and an inverse agonist of ERR α and ERR γ . Kaempferol has a wide range of antitumor, anti-inflammatory, antioxidant, antibacterial and antiviral activities.
Targets(IC50)	Apoptosis, Estrogen Receptor/ERR, Mitophagy, Estrogen/progestogen Receptor, HIV Protease, Endogenous Metabolite, Parasite, Autophagy
In vitro	<p>METHODS: Human breast cancer cells MDA-MB-231 were treated with Kaempferol (0.01-100 μM) for 72 h, and cell viability was detected by MST assay.</p> <p>RESULTS: Kaempferol inhibited cell proliferation in a dose-dependent manner with an IC₅₀ value of 43 μmol/L. [1]</p> <p>METHODS: Human gastric cancer cells AGS and SNU638 were treated with Kaempferol (50 μM) for 8-24 h. The expression levels of target proteins were detected by Western Blot.</p> <p>RESULTS: Kaempferol increased cleaved caspase-3 and -9 and decreased Bcl-2 levels in a time-dependent manner. [2]</p>
In vivo	<p>METHODS: To investigate the effects on hemorrhagic shock mice, Kaempferol (10 mg/kg) was administered intraperitoneally to C57/BL6 mice experiencing hemorrhagic shock.</p> <p>RESULTS: Pre-treatment of hemorrhagic shock mice with Kaempferol significantly reduced plasma TNF-α and IL-6 levels; restored MPO, SOD and MDA in the heart, lungs and liver; and increased HO-1 expression in the same organs. [3]</p> <p>METHODS: To study the anti-neuroinflammatory effects, Kaempferol (25-100 mg/kg) was administered by gavage to BALB/c mice once a day for seven days, and LPS (5 mg/kg) was injected on the seventh day.</p> <p>RESULTS: Kaempferol may be a promising neuroprotective agent to alleviate inflammatory responses and blood-brain barrier dysfunction by inhibiting HMGB1 release and downregulating the TLR4/MyD88 inflammatory pathway. [4]</p>
Kinase Assay	Right atria or sinus nodal cells are homogenized in lysis buffer consisting of (50 mM Tris-HCl pH 7.5, 100 mM KCl, 1 mM ethylenediamine tetraacetic acid, 1 mM ethylene glycol tetraacetic acid, 1 mM dithiothreitol, 0.1 mM phenylmethylsulfonyl fluoride, 0.5 mM Benzimidazole, 20 mg/L Leupeptin, 20 mM sodium pyrophosphate, 50 mM NaF, and 50 mM sodium β -glycerophosphate), and total protein content is determined by the Bradford assay. Caspase-3 activity is determined by EnzChek Caspase-3 Assay Kit[3].
Cell Research	Kaempferol is prepared in DMSO (100 mM) and stored (-20°C), and then diluted with appropriate medium[2]. Ovarian cancer cells are seeded in 96-well plates at 2000

cells/well and incubated overnight before treatment with 0-160 μ M Kaempferol for 24 hours in triplicates. The medium is removed, and the plates are freeze-thawed to lyse cells. Each well is added with 200 μ L 1 \times CyQUANT cell lysis buffer containing 5 \times SYBR Green I and incubated at room temperature (RT) for 5 minutes. The reaction (50 μ L) is transferred to PCR strip tubes and the fluorescent signal is measured at 90°C with a real-time Chromo4 PCR instrument. To ensure that cell proliferation assays are performed within a linear range of cell numbers, a standard curve is generated by seeding different amount of OVCAR-3 cells (based on counting with a hemacytometer) in a 96-well plate, and measuring genomic DNA abundance after overnight incubation. Three independent experiments are performed and data is pooled for statistical analysis[2].

Solubility Information

Solubility	H2O: < 1 mg/mL (insoluble or slightly soluble), Ethanol: 28.6 mg/mL (100 mM), DMSO: 45 mg/mL (157.21 mM), & 1 mg/ml refers to the product slightly soluble or insoluble)
------------	---

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.4936 mL	17.4679 mL	34.9357 mL
5 mM	0.6987 mL	3.4936 mL	6.9871 mL
10 mM	0.3494 mL	1.7468 mL	3.4936 mL
50 mM	0.0699 mL	0.3494 mL	0.6987 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Wang R, Deng Z, Zhu Z, et al. Kaempferol promotes non-small cell lung cancer cell autophagy via restricting Met pathway. *Phytomedicine*. 2023: 155090.
Zhu L, et al. Kaempferol Suppresses Proliferation and Induces Cell

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 36 Washington Street, Wellesley Hills, MA 02481