

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

Data Sheet (Cat.No.T2376)



Liproxstatin-1

Biological Description

Chemical Proper	ties
CAS No. :	950455-15-9
Formula:	C19H21CIN4
Molecular Weight:	340.85
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year

Description Liproxstatin-1 is a potent and selective inhibitor of ferroptosis (IC50=22 nM). Liproxstatin-1 protects cells from ferroptosis induced by ferroptosis inducers (e.g., Erastin, RSL3). Targets(IC50) Ferroptosis In vitro **METHODS**: Mouse fibroblasts L929 were treated with Liproxstatin-1 (0-250 nM) for 24 h. Cell viability was measured using AquaBluer. **RESULTS**: Liproxstatin-1 protected against FINs such as BSO (10 μ M), erastin (1 μ M), and RSL3 (0.5 μ M) in a dose-dependent manner, while it did not rescue staurosporine (0.2 μ M) or H2O2 (200 μ M)-induced cell death. [1] **METHODS**: Oligodendrocytes OLN93 were treated with Liproxstatin-1 (1 μ M) and RSL-3 (7.89 µM) for 24 h, and GSH levels were measured by micro reduced GSH assay kit. **RESULTS**: Liproxstatin-1 treatment increased GSH levels compared to the RSL-3 group. [2] In vivo **METHODS:** To test the potential to prevent animal-induced Gpx4 destruction in vivo, Liproxstatin-1 (10 mg/kg) was administered intraperitoneally to TAM-treated CreERT2; Gpx4fl/fl mice once daily for two weeks. **RESULTS**: Liproxstatin-1 significantly prolonged survival. the number of TUNEL+ cells in the Liproxstatin-1 group was significantly reduced, and the table Liproxstatin-1 delayed the desmoplasia of renal tubular cells. [1] **METHODS**: To investigate the role of ferroptosis in inflammation-associated cognitive deficits, Liproxstatin-1 (10 mg/kg) was administered intraperitoneally to C57BL/6 mice once daily for five days followed by LPS administration.

RESULTS: Liproxstatin-1 ameliorated memory deficits in a mouse model of LPS-induced
cognitive impairment. The protective effects of Liproxstatin-1 were associated with
attenuation of iron deposition and modulation of the iron-death-associated protein
families, TF, xCT, Fth, Gpx4 and FtMt. [3]Cell ResearchCell viability is assessed at different time points after treatment using AquaBluer
according to the manufacturer’s recommendations. Alternatively, cell death is
also quantified by measuring released lactate dehydrogenase activity using the

cytotoxicity detection kit (LDH). (Only for Reference)

Solubility Information

A DRUG SCREENING EXPERT

Solubility	Ethanol: soluble,
	H2O: Insoluble,
	DMSO: 60 mg/ml (176.03 mM)
	(< 1 mg/ml refers to the product slightly soluble or insoluble)

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.9338 mL	14.6692 mL	29.3384 mL
5 mM	0.5868 mL	2.9338 mL	5.8677 mL
10 mM	0.2934 mL	1.4669 mL	2.9338 mL
50 mM	0.0587 mL	0.2934 mL	0.5868 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Hu G, Cui Z, Chen X, et al.Suppressing Mesenchymal Stromal Cell Ferroptosis Via Targeting a Metabolism-Epigenetics Axis Corrects their Poor Retention and Insufficient Healing Benefits in the Injured Liver Milieu.Advanced

Inhibitor • Natural Compounds • Compound Libraries • Recombinant ProteinsThis product is for Research Use Only• Not for Human or Veterinary or Therapeutic UseTel:781-999-4286E_mail:info@targetmol.comAddress:36 Washington Street,Wellesley Hills,MA 02481