



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)


[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Digitonin

## Chemical Properties

CAS No. :	11024-24-1
Formula:	C <sub>56</sub> H <sub>92</sub> O <sub>29</sub>
Molecular Weight:	1229.31
Appearance:	no data available
Storage:	Powder: -20°C for 3 years   In solvent: -80°C for 1 year



## Biological Description

Description	Digitonin is a natural product of the glycoside family and is a detergent. Digitonin has antitumor activity, binds to cholesterol molecules to increase the permeability of cell membranes, and is active in the cleavage of cell membranes in a wide range of cells.
Targets(IC50)	Others
In vitro	<p><b>METHODS:</b> Rat adrenal pheochromocytoma cells PC-12 were treated with Digitonin (10 µg/mL) for 10 min to increase cell permeability.</p> <p><b>RESULTS:</b> After permeabilization with Digitonin, PC12 cells could be stimulated to release norepinephrine by micromolar concentrations of free Ca<sup>2+</sup>. [1]</p> <p><b>METHODS:</b> Human glioblastoma cells U87 were treated with Digitonin (1-5 µg/mL) for 20 min, and cell viability was measured by MTT after overnight cell incubation.</p> <p><b>RESULTS:</b> 1 µg/mL Digitonin treatment resulted in a slight increase in cell viability, 3 µg/mL Digitonin reduced the growth-promoting effect, and the cell viability was similar to that of the control, and 5 µg/mL Digitonin inhibited the cell viability, which indicated a significant toxicity to the cells. [2]</p> <p><b>METHODS:</b> Immortalized human T-lymphocytes Jurkat were treated with Digitonin (1-4 µg/mL) for 20 min and cell viability was measured by Resazurin assay.</p> <p><b>RESULTS:</b> Digitonin treatment resulted in a significant decrease in metabolic activity and this inhibition was concentration-dependent. [2]</p>
In vivo	<p><b>METHODS:</b> To determine whether Digitonin enhances the efficacy of CBDCA in vivo, Digitonin (0.2 mL 400 µM) was hepatic arterially infused into LH rats bearing hepatocellular carcinoma tumors, followed by injection of CBDCA 10 min later.</p> <p><b>RESULTS:</b> Digitonin pretreatment increased the tumor uptake of CBDCA and enhanced the cytotoxic effect of CBDCA. [3]</p>

## Solubility Information

Solubility	DMSO: 27.5 mg/mL (22.37 mM), ( $< 1$ mg/ml refers to the product slightly soluble or insoluble)
------------	--

### Preparing Stock Solutions

---

	1mg	5mg	10mg
1 mM	0.8135 mL	4.0673 mL	8.1346 mL
5 mM	0.1627 mL	0.8135 mL	1.6269 mL
10 mM	0.0813 mL	0.4067 mL	0.8135 mL
50 mM	0.0163 mL	0.0813 mL	0.1627 mL

---

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

### Reference

Schäfer T, et al. Digitonin-permeabilized cells are exocytosis competent. *J Neurochem.* 1987 Dec;49(6):1697-707.  
Chen Y, Wang X, Shi H, et al. Montelukast Inhibits HCoV-OC43 Infection as a Viral Inactivator. *Viruses.* 2022, 14(5):

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481