



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Stattic

Chemical Properties

CAS No. :	19983-44-9
Formula:	C ₈ H ₅ NO ₄ S
Molecular Weight:	211.19
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Stattic (STAT3 Inhibitor V) is a STAT3 inhibitor (IC ₅₀ =5.1 μM) that selectively inhibits STAT3 activation, dimerization, and nuclear translocation. Stattic has antitumor activity and induces apoptosis.
Targets(IC ₅₀)	Apoptosis,STAT
In vitro	<p>METHODS: Human pancreatic cancer cells PANC-1 and BxPc-3 were treated with Stattic (1-10 μM) for 12-48 h. Cell viability was measured using the CCK-8.</p> <p>RESULTS: Stattic decreased the proliferation of PANC-1 and BxPc-3 cells in a concentration- and time-dependent manner, and the IC₅₀s of Stattic on BxPc-3 and PANC-1 cells were 3.135-5.296 μM and 3.835-4.165 μM, respectively, after treatment with Stattic for 24 h. [1]</p> <p>METHODS: Human hepatocellular carcinoma cells HepG2 were treated with Stattic (5-20 μM) for 1 h, followed by stimulation with IL-6 or IFN-γ, and the expression levels of target proteins were detected by Western Blot.</p> <p>RESULTS: Pre-incubation with Stattic resulted in a selective decrease in the phosphorylation of STAT3 Tyr705, while the activation of STAT1 Tyr701 remained unchanged. [2]</p>
In vivo	<p>METHODS: To test the antitumor activity in vivo, Stattic (10 mg/kg) was administered intraperitoneally once daily for four weeks to BALB/c nude mice harboring human pancreatic adenocarcinoma tumor PANC-1.</p> <p>RESULTS: Stattic inhibited PC growth in a nude mouse tumor model by inactivating STAT3. [1]</p> <p>METHODS: To investigate the role in acute liver injury, Stattic (5 mg/kg in DMSO:olive oil = 1:19) was administered as a single intraperitoneal injection to BALB/c mice with LPS/d-GalN induced acute liver injury.</p> <p>RESULTS: Stattic was protective against LPS/d-GalN-induced liver injury, and its protective effect may be related to its anti-inflammatory and anti-apoptotic effects. [3]</p>
Kinase Assay	The screening was performed at approximately 30°C. The specificity of screening hits was validated in analogous assays for binding of the test compounds to the SH2 domains of STAT1, STAT5, and Lck. The final concentration of buffer components used for all FP assays was 10 mM HEPES (pH 7.5), 1 mM EDTA, 0.1% Nonidet P-40, 50 mM NaCl, and 10% DMSO. The absence of dithiothreitol is essential for inhibitory activity. The sequences of the peptides were: STAT3, 5-carboxyfluorescein-GY(PO ₃ H ₂)LPQTV-NH ₂ ; STAT1, 5-carboxyfluorescein-GY(PO ₃ H ₂)DKPHVL; STAT5, 5-carboxyfluorescein-GY (PO ₃ H ₂)

LVLDKW; and Lck, 5-carboxyfluorescein-GY(PO3H2)EEIP. Peptides were >95% pure. For specificity analysis at 30°C, proteins were used at 150 nM (STAT1, STAT3, and STAT5). For specificity analysis at 37°C, proteins were used at 370 nM (STAT3) or 100 nM (Lck). Proteins were incubated with test compounds in tubes at the indicated temperatures for 60 min prior addition of the respective 5-carboxyfluorescein labeled peptides (final concentration: 10 nM). Analysis of c-Myc/Max and Jun/Jun dimerization and DNA binding at 37°C was performed as described but in the absence of DTT. Before measurement at room temperature, the mixtures were allowed to equilibrate for at least 30 min. Test compounds were used at the indicated concentrations diluted from 20× stock in DMSO. Binding curves and inhibition curves were fitted with SigmaPlot. All competition curves were repeated three times in independent experiments. For the analysis of time dependence of the inhibition, the components were mixed from stock solutions kept at 0°C and then incubated at 37°C. Aliquots were taken at the indicated time points [1].

Cell Research

MDA-MB-231, MDA-MB-435S, and MDA-MB-453 cells were seeded. at 5×10^4 cells in 6-well plates, grown for 24 hr before adding DMSO or Stattic (final DMSO concentration 0.1%) and then incubated with the inhibitor for 24 hr. All cells were collected and resuspended in buffer (0.1% sodium citrate, 0.1% Triton X-100, 20 μ M propidium iodide) and incubated for 3 hr before 10^4 cells per sample were analyzed by flow cytometry with a FACSCalibur equipped with a 488 nm laser [1].

Solubility Information

Solubility

Ethanol: 1.1 mg/mL (5 mM),
DMSO: 55 mg/mL (260.43 mM),
(< 1 mg/ml refers to the product slightly soluble or insoluble)

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	4.7351 mL	23.6754 mL	47.3507 mL
5 mM	0.947 mL	4.7351 mL	9.4701 mL
10 mM	0.4735 mL	2.3675 mL	4.7351 mL
50 mM	0.0947 mL	0.4735 mL	0.947 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Guo H, et al. Inhibition of STAT3Y705 phosphorylation by Stattic suppresses proliferation and induces mitochondrial-dependent apoptosis in pancreatic cancer cells. Cell Death Discov. 2022 Mar 14;8(1):116.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 36 Washington Street, Wellesley Hills, MA 02481