

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



# Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

# Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# Data Sheet (Cat.No.T6834)



#### **FCCP**

#### **Chemical Properties**


CAS No.: 370-86-5

Formula: C10H5F3N4O

Molecular Weight: 254.17

Appearance: no data available

Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year



#### **Biological Description**

Description	FCCP (Trifluoromethoxy carbonylcyanide phenylhydrazone) is an oxidative phosphorylation (OXPHOS) inhibitor and mitochondrial proton carrier uncoupler. often used as an apoptosis inducer.				
Targets(IC50)	ATPase,Mitochondrial Metabolism				
In vitro	METHODS: Rat ventricular myocytes were treated with FCCP (10-1000 nM) for 1400 sec, and cellular oxygen consumption was measured using a Clark Oxygen Electrode.  RESULTS: Oxygen consumption increased significantly, immediately, and in a dose-dependent manner after the addition of FCCP. [1]  METHODS: Rat adrenal medullary chromaffinoma cells PC12 were treated with FCCP (30 μM) for 0.5-2 h. The rate of protein synthesis was measured using [3H]methionine.  RESULTS: FCCP treatment produced a strong inhibition (68%) of protein synthesis rate for at least 2 hours. [2]				
In vivo	METHODS: To test the effect on stroke, FCCP (1 mg/kg) was injected intraperitoneally into a C57BL/6J mouse model of stroke, followed by one hour of transient middle cerebral artery occlusion (tMCAO).  RESULTS: Infarct volumes in the cortex, striatum, and whole hemisphere were significantly increased in mice pretreated with FCCP. Mice receiving FCCP had significantly increased neurologic deficit scores compared to carriers. [3]  METHODS: To assay anti-tumor activity in vivo, FCCP (1 mg/kg) and cisplatin (2 mg/kg) were intraperitoneally injected every two days for two weeks into C57BL/6 mice harboring mouse ovarian epithelial carcinoma tumor ID8.  RESULTS: The combination of FCCP and cisplatin inhibited tumor growth via OMA1-induced mitochondrial and ER stress. [4]				
Cell Research	Protein synthesis rate is assayed in 24-mm diameter multi-well dishes with fresh medium containing 0.175 Ci/mmol of [3H]methionine (200 μ M), for 30 min at 37° C. PC12 cells are treated with FCCP for different period of times. (Only for Reference)				

## **Solubility Information**

Solubility	DMSO: 55 mg/mL (216.39 mM),	
	(< 1 mg/ml refers to the product slightly soluble or insoluble)	

Page 1 of 2 www.targetmol.com

#### **Preparing Stock Solutions**

	1mg	5mg	10mg
1 mM	3.9344 mL	19.6719 mL	39.3437 mL
5 mM	0.7869 mL	3.9344 mL	7.8687 mL
10 mM	0.3934 mL	1.9672 mL	3.9344 mL
50 mM	0.0787 mL	0.3934 mL	0.7869 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

#### Reference

Brennan JP, et al. FCCP is cardioprotective at concentrations that cause mitochondrial oxidation without detectable depolarisation. Cardiovasc Res. 2006 Nov 1;72(2):322-30.<a href="https://br/>Liu H, Sun Y, Xu H, et al.">br/>Liu H, Sun Y, Xu H, et al.</a> PTEN-induced putative

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481

Page 2 of 2 www.targetmol.com