



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Glucoraphanin

Chemical Properties

CAS No. :	21414-41-5
Formula:	C ₁₂ H ₂₃ NO ₁₀ S ₃
Molecular Weight:	437.51
Appearance:	no data available
Storage:	store at low temperature,store under nitrogen,keep away from direct sunlight,keep away from moisture Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Glucoraphanin, the precursor of the widely extolled chemopreventive agent sulforaphane found in broccoli, has antioxidant activity.
Targets(IC50)	Antioxidant,Reactive Oxygen Species
In vivo	Glucoraphanin reduces weight gain, enhances energy expenditure, improves insulin sensitivity, and glucose tolerance in HFD-fed mice, but it does not exert these effects in Nrf2 ^{-/-} . It blocks HFD-induced reduction of Ucp1 protein levels in white adipose depots of wild-type mice but not in Nrf2 ^{-/-} , alleviates HFD-induced hepatic steatosis and oxidative stress, and suppresses HFD-induced proinflammatory activation of macrophages in the liver and adipose tissue [1]. Additionally, pellets containing 0.1% Glucoraphanin (GF) significantly attenuate decreased social avoidance time and sucrose preference in stressed mice [2].
Animal Research	Male C57BL/6JSlc mice are 7 weeks of age. The Nrf2 knockout (Nrf2 ^{0/0}) mouse strain is used in the assay. After 1 week of acclimation, mice are fed normal chow (NC) (containing 2.2% dextrinized cornstarch, 10% kcal from fat), NC containing 0.3% glucoraphanin (NC-GR) (containing 2.2% extract powder), an HFD (containing 2.2% dextrinized cornstarch, 60% kcal from fat), or an HFD containing 0.3% glucoraphanin (HFD-GR) (containing 2.2% extract powder) for 14 weeks. Both the NC and the HFD containing cornstarch or glucoraphanin are prepared. All mice studied are maintained on a 12-h light/dark cycle at 24-26°C with free access to water and food [1].

Solubility Information

Solubility	H ₂ O: 74 mg/mL (169.14 mM), (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.2857 mL	11.4283 mL	22.8566 mL
5 mM	0.4571 mL	2.2857 mL	4.5713 mL
10 mM	0.2286 mL	1.1428 mL	2.2857 mL
50 mM	0.0457 mL	0.2286 mL	0.4571 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Sun R, Huang H, Wang Z, et al. Computer-driven Evolution of Myrosinase from the Cabbage Aphid for Efficient Production of (R)-Sulforaphane. *Journal of Agricultural and Food Chemistry*. 2024

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 36 Washington Street, Wellesley Hills, MA 02481