



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Solanum Tuberosum (Potato) Lectin (STL, PL), Biotinylated

B-1165-2

[Product Images](#)



Short Description

This lectin binds oligomers of *N*-acetylglucosamine and some bacterial cell wall oligosaccharides containing *N*-acetylglucosamine and *N*-acetylmuramic acid. Although the carbohydrate binding specificity is similar to wheat germ agglutinin and *Datura stramonium* lectin, several differences have been reported for potato lectin.

Biotinylated *Solanum tuberosum* lectin has an appropriate number of biotins bound to provide the optimum staining characteristics for this lectin. This conjugate is supplied essentially free of unconjugated biotins and is preserved with sodium azide.

Additional Information

Unit Size	2 mg
Applications	Immunohistochemistry / Immunocytochemistry, Immunofluorescence, Blotting Applications, Elispot, ELISAs, Glycobiology
Recommended Usage	For most applications we recommend a freshly prepared working solution of 5-20 µg/ml in the above buffer.
Recommended Storage	2-8 °C; Store frozen for long term storage
Solution	10 mM HEPES, pH 7.5, 0.15 M NaCl, 0.08% sodium azide, 0.1 mM CaCl ₂ , 20 mM N-acetylglucosamine.
Concentration	2 mg active conjugate/ml
Conjugate	Biotinylated
Sugar Specificity	N-Acetylglucosamine

