



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Griffonia (Bandeiraea) Simplicifolia Lectin II (GSL II, BSL II), Biotinylated B-1215-2

[Product Images](#)



Short Description

Griffonia (Bandeiraea) Simplicifolia Lectin II is a dimeric glycoprotein composed of two subunits of nearly identical size with each subunit having disulfide-linked chains and a binding site for α - or β -linked *N*-acetylglucosamine residues. Unlike other *N*-acetylglucosamine specific lectins, increasing the number of *N*-acetylglucosamine residues beyond two does not improve affinity. GSL II has been reported to be unique in its ability to recognize exclusively α - or β -linked *N*-acetylglucosamine residues on the nonreducing terminal of oligosaccharides.

Biotinylated GSL II has an appropriate number of biotins bound to provide the optimum staining characteristics for this lectin. This conjugate is supplied essentially free of unconjugated biotins and is preserved with sodium azide.

Additional Information

| | |
|---------------------|---|
| Unit Size | 2 mg |
| Applications | Immunohistochemistry / Immunocytochemistry, Immunofluorescence, Blotting Applications, Elispot, ELISAs, Glycobiology |
| Recommended Usage | Reconstitute biotinylated lectin by addition 1 ml of water. The resulting solution will have the following composition: 10 mM HEPES, pH 7.5, 0.15 M NaCl, 0.08% sodium azide, 0.1 mM CaCl ₂ , 10 mM <i>N</i> -Acetylglucosamine. For most applications, we recommend a freshly prepared working solution of 5-20 μ g/ml in the above buffer. |
| Recommended Storage | 2-8 °C; Store frozen for long term storage |
| Conjugate | Biotinylated |
| Sugar Specificity | <i>N</i> -Acetylglucosamine |

