



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Wisteria Floribunda Lectin (WFA, WFL), Biotinylated

B-1355-2

[Product Images](#)



Short Description

The binding specificity of *Wisteria floribunda* lectin (WFL) is not completely clear but this lectin appears to preferentially bind carbohydrate structures terminating in *N*-acetylgalactosamine linked α or β to the 3 or 6 position of galactose. This lectin has been used to fractionate lymphocyte populations, and although not mitogenic, elicits the production of lymphokines from murine splenocytes.

Biotinylated WFL has an appropriate number of biotins bound to provide the optimum staining characteristics for this lectin. This conjugate is supplied essentially free of unconjugated biotins and is preserved with sodium azide.

Additional Information

Unit Size	2 mg
Applications	Immunohistochemistry / Immunocytochemistry, Immunofluorescence, Blotting Applications, Elispot, ELISAs, Glycobiology
Recommended Usage	For most applications, we recommend a freshly prepared working solution of 5-20 $\mu\text{g/ml}$ in the above buffer.
Recommended Storage	2-8 °C; Store frozen for long term storage
Solution	10 mM HEPES, pH 7.5, 0.15 M NaCl, 0.08% sodium azide, 0.1 mM CaCl_2 .
Concentration	2 mg active conjugate/ml
Conjugate	Biotinylated
Sugar Specificity	N-Acetylgalactosamine

