



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Goat Anti-Rat IgG Antibody, mouse adsorbed (H+L), Biotinylated

BA-9401-.5

[Product Images](#)



Short Description

Biotinylated Goat Anti-Rat IgG, mouse adsorbed, secondary antibody is used in an avidin-biotin or streptavidin-biotin detection system. This adsorbed antibody allows primary antibodies made in rat to be used on mouse tissues.

Features:

- Recognizes both heavy and light chains (H+L)
- Biotinylated to ensure the maximum degree of labeling without compromising the specificity or affinity of the antibody
- Can be used for tissue and cell staining, ELISAs, and blots
- Supplied in solution
- **The mouse adsorbed anti-rat IgG (H+L) products are designed for use with mouse tissues that may contain endogenous mouse immunoglobulins.**

Additional Information

| | |
|---------------------|---|
| Unit Size | 0.5 mg |
| Applications | Immunohistochemistry / Immunocytochemistry, Immunofluorescence, In situ hybridization, Blotting Applications, Elispot, ELISAs |
| Concentration | 0.5 mg active conjugate/ml |
| Recommended Storage | 2-8 °C; Store frozen for long term storage |
| Solution | 10 mM sodium phosphate, pH 7.8, 0.15 M NaCl, 0.08% sodium azide, 1 mg/ml bovine serum albumin. |
| Recommended Usage | The recommended concentration range for use is 2-10 µg/ml. If this biotinylated antibody is to be used in tissues, which may contain cross-reacting endogenous immunoglobulins, dilution of this biotinylated antibody may be made in buffers containing 2% normal serum from the same species as the tissue. |
| Target Species | Rat |
| Conjugate | Biotinylated |
| Host Species | Goat |
| Format | Concentrate |

