



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Horse Anti-Goat IgG Antibody (H+L), Biotinylated, R.T.U.

BP-9500-50

[Product Images](#)



Short Description

The Ready-to-Use (R.T.U.) Biotinylated Horse Anti-Goat IgG Antibody is used in an avidin-biotin or streptavidin-biotin detection system. These antibodies can be used for tissue and cell staining, ELISAs, and blots.

Features:

- Suitable for use with goat, sheep, and bovine IgG primary antibodies
- Recognizes both heavy and light chains (H+L)

Additional Information

Unit Size	50 ml
Applications	Immunohistochemistry / Immunocytochemistry, Immunofluorescence, In situ hybridization, Blotting Applications, Elispot, ELISAs
Concentration	Ready-to-use solution of biotinylated horse anti-goat IgG in PBS* containing carrier protein and 0.08% sodium azide as preservative.
Recommended Storage	2-8 °C
Recommended Usage	Following incubation with primary antibody, wash specimens in buffer. Apply RTU Biotinylated Anti-Goat IgG and incubate for 30 minutes at room temperature. Wash in buffer for 5 minutes and follow with avidin- or streptavidin-based detection reagent such as RTU VECTASTAIN® Elite® ABC Reagent (PK-7100). * 10mM sodium phosphate, pH 7.8, 0.15M NaCl.
Target Species	Goat
Conjugate	Biotinylated
Host Species	Horse
Format	Ready-to-Use

