



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



---

# Goat Anti-Human IgG Antibody (H+L), Fluorescein

## FI-3000-1.5

[Product Images](#)

---



## Short Description

---

Fluorescein Goat Anti-Human IgG Antibody can be used for immunofluorescence and other applications. Optimal F/P ratios have been established for each conjugate to ensure maximum fluorescence with minimal background staining.

### Features:

- Affinity-purified, ultrapure, high affinity antibody
- Thoroughly adsorbed against serum and immunoglobulins from potentially interfering species
- Recognizes both heavy and light chains (H+L)
- Optimally labeled with fluorescein to provide the brightest label for fluorescence microscopy
- Supplied in solution
- Excitation: 495 nm
- Emission: 515 nm
- Color: Green

## Additional Information

Unit Size	1.5 mg
Applications	Immunofluorescence, Blotting Applications, Flow Cytometry/Cell Separation
Concentration	1.5 mg active conjugate/ml
Recommended Storage	2-8 °C
Solution	10 mM HEPES, 0.15 M NaCl, pH 7.5, 0.08% sodium azide.
Maximum Emission	510-520 nm
Maximum Excitation	490-500 nm
Recommended Usage	The recommended concentration range for use is 5-20 µg/ml. If this fluorescein-labeled antibody is to be used in tissues which may contain cross-reacting endogenous immunoglobulins, dilution of this antibody may be made in buffers containing 2% normal serum from the same species as the tissue.
Target Species	Human
Conjugate	Fluorescein
Color of Fluorescence	Green
Host Species	Goat
Format	Concentrate

