



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Horse Anti-Goat IgG Antibody (H+L), Peroxidase

PI-9500-1

[Product Images](#)



Short Description

Features:

- Affinity-purified, ultrapure, high affinity antibody
- Suitable for use with goat, sheep, and bovine IgG primary antibodies
- Thoroughly adsorbed against serum and immunoglobulins from potentially interfering species
- Unless otherwise specified, antibodies recognizes both heavy and light chains (H+L)
- Cross-linked with the highest specific activity horseradish peroxidase
- Supplied in solution

Additional Information

Unit Size	1 mg
Applications	Immunohistochemistry / Immunocytochemistry, In situ hybridization, Blotting Applications, Elispot, ELISAs
Concentration	1 mg active conjugate/ml
Recommended Storage	2-8 °C, DO NOT FREEZE
Solution	10 mM HEPES, 0.15 M NaCl, pH 7.5, preservative
Recommended Usage	<p>Important Note: The use of bovine products such as casein, serum, albumin or non-fat dry milk as blocking agents in ELISA or blot assays may produce a high background due to cross-reactivity of anti-goat IgG with bovine immunoglobulins that may be present. If this antibody is to be used in these applications, blocking with Animal-Free blocker™ solution (Cat. No. SP-5030) is recommended. Recommended diluent for microtitration plate assays: 10 mM phosphate, 0.15 M NaCl, pH 7.5, 0.1% Tween 20. Working concentration of HRP-anti goat IgG is 0.5-5.0 µg/ml.</p>
Target Species	Goat
Conjugate	Peroxidase
Host Species	Horse
Format	Concentrate
Safety Title	Cancer

