



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Griffonia (Bandeiraea) Simplicifolia Lectin I (GSL I, BSL I), Rhodamine

RL-1102-2

[Product Images](#)



Short Description

GSL I is a family of glycoproteins with molecular weights of approximately 114 kDa. There are two types of subunits, termed "A" and "B", with slightly different molecular weights. These subunits combine to form tetrameric structures, resulting in five isolectins. The "A"-rich lectin preferentially agglutinates blood group A erythrocytes and thus appears to be specific for α -N-acetylgalactosamine residues, while the "B"-rich lectin preferentially agglutinates blood group B cells and is specific for α -galactose residues. Our GSL I is a mixture of the five isolectins. GSL I has been reported to bind several glycoproteins including laminin.

Rhodamine labeled GSL I has an appropriate number of fluorochromes bound to provide the optimum staining characteristics for this lectin. This conjugate is supplied essentially free of unconjugated fluorochromes. The excitation maximum is at 550 nm and the emission maximum is at 575 nm.

Additional Information

Unit Size	2 mg
Applications	Immunofluorescence, Glycobiology
Recommended Usage	The recommended concentration range for use is 5-20 μ g/ml.
Recommended Storage	2-8 °C
Maximum Excitation	545-555 nm
Maximum Emission	570-580 nm
Solution	10 mM HEPES, 0.15 M NaCl, pH 7.5, 0.08% sodium azide, 0.1 mM CaCl ₂
Concentration	2 mg active conjugate/ml
Conjugate	Rhodamine
Color of Fluorescence	Red
Sugar Specificity	Galactose, N-Acetylgalactosamine

