

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in







## Oligo Clean & Concentrator™

Clean-up DNA/RNA oligos from any reaction

#### **Highlights**

- Quick, 2-minute clean-up of ultra-pure DNA RNA and oligonucleotides.
- Complete removal of dyes, salts, enzymes, nucleotides and short oligos.
- Eluted DNA/RNA (≥ 6 µl) is ready for hybridization, sequencing, PCR, ligation, etc.

Catalog Numbers: D4060, D4061



Scan with your smart-phone camera to view the online protocol/video.







## **Table of Contents**

Product Contents	01
Specifications	02
Applications	02
Product Description	03
Protocol	04
(I) Buffer Preparation	04
(II) DNA/RNA Clean-Up	05
Appendices	06
First-Strand cDNA Clean-Up (following RT)	06
Format Compatibility	06
Ordering Information	07
Complete Your Workflow	08
Guarantee	09

Revised on: 2/16/2023

#### **Product Contents**

Oligo Clean & Concentrator™	<b>D4060</b> (50 prep)	<b>D4061</b> (200 prep)
Oligo Binding Buffer	10 ml	40 ml
DNA Wash Buffer (concentrate) <sup>1</sup>	24 ml	48 ml
Zymo-Spin <sup>™</sup> IC Columns	50	200
Collection Tubes	100	400
Instruction Manual	1	1

**Storage Temperature** - Store all kit components (i.e., buffers, columns) at room temperature. Before use:

<sup>1</sup> Prior to use, add 96 ml 100% ethanol (104 ml 95% ethanol) to the 24 ml **DNA Wash Buffer** concentrate (D4060) or 192 ml 100% ethanol (208 ml of 95% ethanol) to the 48 ml **DNA Wash Buffer** concentrate (D4061).

### **Specifications**

- Sample Sources Enzymatic reactions mixtures containing oligonucleotides ≥ 16 nt (radioactive-, biotin-, DIG-labeled, etc.)
- Size Limits For oligonucleotides ≥ 16 nt, up to 23 kb.
- Purity A<sub>260</sub>/A<sub>280</sub> & A<sub>260</sub>/A<sub>230</sub> > 1.8. DNA/RNA is ready for hybridization, sequencing, ligation, PCR and etc.
- Binding Capacity 10 µg ssDNA/RNA or 5 µg dsDNA with a typical recovery of > 90% (Zymo-Spin<sup>™</sup> IC Column).
- Elution Volume ≥ 6 µl (water not provided).
- Detergent Tolerance ≤ 5% Triton X-100, ≤ 5% Tween-20, ≤ 5% Sarkosyl, ≤ 0.1% SDS.
- Equipment Needed (user provided) Microcentrifuge.


#### **Applications**


- Isotope and Dye Removal Efficiently removes unincorporated fluorescent (i.e., AMCA, FITC, BIO, DIG, Cy3, Cy5, FAM, etc.) and radiolabeled dNTP derivatives from DNA following in vitro labeling reactions.
- DNA Fragment Clean-up from Enzymatic Reactions Desalting of DNA with the removal of DNA polymerases, modifying enzymes, RNA polymerases, ligases, kinases, nucleases, phosphatases, endonucleases, etc.
- Post-Reverse Transcription (RT) and cDNA Clean-up –
   Purifies DNA following RT, either as a DNA/RNA complex or as single stranded cDNA following chemical hydrolysis of the RNA template (page 6).

### **Product Description**

The **Oligo Clean & Concentrator**™ kit provides a streamlined method for efficient recovery and clean-up of DNA and RNA oligonucleotides ≥16 nt from labeling (radioactive, biotin, DIG, *etc.*) and other enzymatic reactions. Unincorporated nucleotides, short oligos, dyes, enzymes, and salts are effectively removed by the clean-up procedure.

There is no need for organic denaturants or chloroform. Instead, the kit features Zymo- $Spin^{\mathbb{M}}$  column technology and employs a single-buffer system that allows for efficient oligonucleotide adsorption to the matrix of Zymo- $Spin^{\mathbb{M}}$  IC Column. Oligonucleotide is washed and concentrated into a small volume of water ( $\geq$  6  $\mu$ I). Purified oligonucleotide, available in just 2 minutes, is suitable for hybridization, gel shift assays, enzymatic reactions, ligation, sequencing, microarray analysis, *etc*.





Oligo Clean & Concentrator<sup>™</sup> facilitates > 90% recovery of ssDNA oligonucleotides (A) and efficient short oligo and nucleotide removal (B).

#### **Protocol**

The protocol consists of: (I) Buffer Preparation and (II) DNA/RNA Clean-Up

#### (I) Buffer Preparation

✓ Prior to use, add 96 ml 100% ethanol (104 ml 95% ethanol) to the 24 ml DNA Wash Buffer concentrate (D4060) or 192 ml 100% ethanol (208 ml of 95% ethanol) to the 48 ml DNA Wash Buffer concentrate (D4061).

#### (II) DNA/RNA Clean-up

- ✓ Perform all steps at room temperature and centrifugation at 10,000-16,000 x g for 30 seconds, unless specified.
- ✓ Scale up the volumes proportionally (steps 1-2), if needed.
- Add 100 μl Oligo Binding Buffer to 50 μl sample¹.
- 2. Add 400 µl ethanol<sup>2</sup> (95-100%) and mix well by pipetting.
- 3. Transfer the sample to the **Zymo-Spin**<sup>™</sup> **IC Column**<sup>3</sup> in a **Collection Tube** and centrifuge. Discard the flow-through<sup>4</sup>.
- 4. Add 750 μl **DNA Wash Buffer** to the column and centrifuge for 1 minute ensure complete removal of the wash buffer. Carefully, transfer the column into a nuclease-free tube (not provided).
- 5. Add 15 μl water<sup>5</sup> directly to the column matrix and centrifuge.

Alternatively, for highly concentrated DNA/RNA use  $\geq$  6  $\mu$ l elution.

The eluted oligonucleotide can be used immediately or stored frozen.

<sup>1</sup> To minimize pipetting error, adjust the sample volume to 50  $\mu$ l (minimum).

<sup>2</sup> For DNA/RNA ≥ 80 nt, only 200 µl ethanol is required.

<sup>3</sup> To process samples >700 µl, **Zymo-Spin**™ columns may be reloaded.

<sup>4</sup> For radioactive samples, transfer the column into a new Collection Tube and discard the tube containing the radioactive flow-through appropriately.

<sup>5</sup> Alternatively, TE buffer can be used for elution (if required).

### **Appendices**

#### cDNA Clean-Up following Reverse Transcription (RT)

The Oligo Clean & Concentrator can be used to effectively clean and concentrate first-strand cDNA following reverse transcription (RT) and hydrolysis. The Oligo Binding Buffer will neutralize the hydrolysis reaction and the recovered cDNA may be used directly for microarray analysis. etc.

**Hydrolysis Reaction**: To each 30-50  $\mu$ l RT reaction, add 10  $\mu$ l 0.5 M EDTA and 10  $\mu$ l 1 M NaOH. Then mix and incubate at 65°C for 15 minutes. Proceed to the DNA/RNA Clean-Up protocol, page 5.

### **Format Compatibility**

To adjust binding capacity, simply replace the provided columns with the formats indicated below and follow their respective elution volumes.

Format	Zymo-Spin™ I & IC	Zymo-Spin™ II & IICR	Zymo-Spin™ III & IIICG	Zymo-Spin™ I-96 Plate
Item Image		V		
Kit Name	OCC™	N/A	N/A	ZR-96 OCC™
Capacity	10 μg ssDNA/RNA or 5 μg dsDNA / prep.	50 μg ssDNA/RNA or 25 μg dsDNA / prep.	100 μg ssDNA/RNA or 50 μg dsDNA / prep.	10 μg ssDNA/RNA or 5 μg dsDNA / prep.
Elution Vol.	≥ 6 µl	≥ 25 µl	≥ 50 µl	≥ 10 µl
Column Cat. Nos.	C1003-50, C1004-50	C1008-50, C1078-50	C1005-50, C1006-50-G	<u>C2004</u>

# **Ordering Information**

Product Description	Catalog No.	Size
Oligo Clean & Concentrator™	D4060 D4061	50 preps. 200 preps.

Individual Kit Components	Catalog No.	Amount
Oligo Binding Buffer	D4060-1-10 D4060-1-40	10 ml 40 ml
DNA Wash Buffer (concentrate)	D4003-2-24 D4003-2-48	24 ml 48 ml
Zymo-Spin™ IC Columns	C1004-50 C1004-250	50 250
Collection Tubes	C1001-50 C1001-500	50 500

## **Complete Your Workflow**

✓ For tough-to-lyse samples in TRIzol, use ZR BashingBead Lysis Tubes:

ZR BashingBead Lysis Tubes	
2.0 mm beads #S6003	For plant/animal tissue
0.1 + 0.5 mm beads #S6012	For microbes
0.1 + 2.0 mm beads #S6014	For microbes in tissue/insects

✓ The only direct, high-throughput and automatable RNA purification from sample lysates in TRIzol (DNase I Set included with all formats):



Direct-zol RNA kits	
Microprep #R2060-R2063	From 1 cell and up
Miniprep #R2050-R2053	Up to 50 ug RNA
Miniprep Plus #R2070-R2073	Up to 100 ug RNA
96-well #R2054-R2057	Spin-plate
MagBeads #R2100-R2105	Automatable (Tecan, Hamilton, Kingfisher, etc.)

✓ For RNA clean-up (purification) from the aqueous phase (e.g., TRIzol, TRI Reagent or similar) or from any enzymatic reaction (e.g., DNase I treated RNA):



RNA Clean & Concentrator kit	
#R1013-R1014	DNase I Set included

✓ For NGS:

Zymo-Seq RiboFree Total RNA Lil	brary Prep kit
#R3000	12 preps
#R3003	96 preps



# 100% satisfaction guarantee on all Zymo Research products, or your money back.

Zymo Research is committed to simplifying your research with quality products and services. If you are dissatisfied with this product for any reason, please call 1(888) 882-9682.

Integrity of kit components is guaranteed for up to one year from date of purchase.

Reagents are routinely tested on a lot-to-lot basis to ensure they provide the highest performance and reliability.

This product is for research use only and should only be used by trained professionals. It is not for use in diagnostic procedures. Some reagents included with this kit are irritants. Wear protective gloves and eye protection. Follow the safety guidelines and rules enacted by your research institution or facility.

Oligo Clean & Concentrator<sup>®</sup> is a registered trademark of Zymo Research Corporation. Other trademarks: TRI Reagent<sup>®</sup>, TRIzol<sup>®</sup> and RNAzol<sup>®</sup> (Molecular Research Center, Inc.), QIAzol<sup>®</sup> (Qiagen GmbH), TriPure<sup>™</sup> (Roche, Inc.), TriSure <sup>™</sup> (Bioline Ltd.), RNAlater<sup>®</sup> (Ambion, Inc.), Bioanalyzer (Agilent Technologies, Inc.).

<sup>™</sup> Trademarks of Zymo Research Corporation

The <b>BEAUTY</b> of <b>SCIENCE</b> is to Make Things <b>SIMPLE</b> ®