



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



ZYMO RESEARCH



Pinpoint™ Slide RNA Isolation System I

RNA from any glass slide tissue section

Highlights

- Spin-column purification of total RNA from tissue mounted on glass slides.
- Pinpoint™ tissue sampling technology is combined with a one-step RNA extraction method.
- RNA is ready for Next-Gen Sequencing, RT/qPCR, etc.

Catalog Numbers:
R1003



Scan with your smart-phone camera to
view the online protocol/video.



tech@zymoresearch.com



www.zymoresearch.com



Toll Free: (888) 882-9682

Table of Contents

Product Contents	01
Specifications	02
Product Description	03
Protocol	04
(I) Buffer Preparation.....	04
(II) Sample Preparation	05
Preparation of Tissue Sections	05
Pinpoint Fractionation	05
(III) Total RNA Purification.....	06
Ordering Information	07
Complete Your Workflow	08
Troubleshooting Guide	09
Notes	10
Guarantee	13

Product Contents

Pinpoint™ Slide RNA Isolation System I	R1003 (50 prep)
Pinpoint™ Solution	1 ml
RNA Extraction Buffer	12 ml
RNA Wash Buffer ¹ (concentrate)	6 ml
DNase/RNase-Free Water	1 ml
Zymo-Spin™ IC Columns	50
Collection Tubes	50
Instruction Manual	1

Storage Temperature - Store all kit components (i.e., buffers, columns) at room temperature.

Before use:

1 Add 24 ml 100% ethanol (26 ml 95% ethanol) to the 6 ml **RNA Wash Buffer** concentrate.



Specifications

- **Sample Sources** – Cells from fresh or frozen tissue sections fixed glass slides¹ by ethanol, acetone, methanol, etc. Use an area of 1 to 100 mm² fresh tissue with 10 µm thickness (approximately 500-1000 cells, depending on the tissue type and cell density).
- **Size** – Total RNA including small/microRNAs (≥ 17 nt).
- **Purity** – A_{260}/A_{280} & $A_{260}/A_{230} > 1.8$. RNA is ready for Next-Gen Sequencing, RT/qPCR, etc.
- **Binding Capacity** – Zymo-Spin™ IC Column yield up to 10 µg RNA.
- **Elution Volume** – ≥ 6 µl DNase/RNase-Free Water.
- **Equipment Needed (user provided)** – Microcentrifuge, vortex, incubator, autoclave.

¹ For paraffin embedded tissue sections, use the Pinpoint™ Slide RNA Isolation System II (R1007).

Product Description

The **Pinpoint™ Slide RNA Isolation System I** is an innovative product designed to isolate RNA from any targeted area of a tissue on microscopic slides. The system combines a powerful **Pinpoint™** tissue sampling method with a unique, single-step RNA extraction/binding buffer that includes **Zymo-Spin™** column technology to yield high quality purified RNA. **Pinpoint™ Slide RNA Isolation System I** allows for the efficient recovery of RNA from fresh tissue sections for any subsequent RNA analyses including RT/qPCR.



RT/qPCR of human tissue section RNA recovered from tissue using the **Pinpoint™ RNA Isolation System I**. Duplicate samples are PCR products from A) human β -actin transcript B) an arbitrary human chromosome 3 transcript.

Protocol

The protocol consists of: (I) Buffer Preparation, (II) Sample Preparation and (III) Total RNA Purification.

(I) Buffer Preparation

- ✓ Add 24 ml 100% ethanol (26 ml 95% ethanol) to the 6 ml **RNA Wash Buffer** concentrate.

(II) Sample Preparation

Preparation of Tissue Sections

1. Clean glass sample slides with sterile ethanol/water and dry by autoclaving (or baking) at 300°C for 4 hours.
2. Mount tissue section ($\geq 10 \mu\text{m}$ thick) onto a glass slide and dry it at 60°C for 30 minutes.
3. To fix the section, submerge the slide in 95% ethanol for 60 minutes.
4. Air dry the sample on the slide for approximately 30 minutes. Proceed to Pinpoint Fractionation, below.

Pinpoint Fractionation (to remove a selected area of tissue from a glass slide)

1. Using a sterile pipette tip, apply 0.5 μl of **Pinpoint™ Solution**¹ per mm^2 of tissue on the slide and gently spread the thick solution over the selected tissue region¹.
2. Allow the **Pinpoint™ Solution** to dry completely as a blue film at room temperature (typically 30-45 minutes), embedding the tissue and cells underneath.
3. Using a sterile blade or scalpel, cut and remove the embedded tissue section from the slide. Then transfer into a nuclease-free tube (not provided).
4. Centrifuge briefly to locate the tissue sample to the bottom of the tube. Proceed to Total RNA Purification, page 6.

¹ Use an area of 1 to 100 mm^2 fresh tissue with 10 μm thickness (approximately 500-1000 cells, depending on the tissue type and cell density).

(III) Total RNA Purification

- ✓ Perform all steps at room temperature and centrifugation at 10,000-16,000 x g for 1 minute, unless specified.
- 1. Add 200 µl **RNA Extraction Buffer** to lyse the tissue sample and pipette up/down several times. Then vortex briefly.
- 2. Incubate the sample on ice for 30 minutes, vortexing briefly every 10 minutes.
- 3. Add 200 µl ethanol (100%) to the sample, mix and then incubate on ice for 10 minutes.
- 4. Transfer the mixture into a **Zymo-Spin™ IC Column¹** in a **Collection Tube** and centrifuge. Discard the flow-through.
- 5. Add 200 µl **RNA Wash Buffer** and centrifuge the column to ensure complete removal of the wash buffer. Then carefully, transfer the column into a nuclease-free tube (not provided).
- 6. Add 10 µl **DNase/RNase-Free Water²** directly to the column matrix and centrifuge.

Alternatively, for highly concentrated RNA use ≥ 6 µl elution.

The eluted RNA can be used immediately or stored frozen.

¹ To process samples > 700 µl, columns may be reloaded.

² To maximize RNA yield, increase the elution volume and/or repeat the elution.

Ordering Information

Product Description	Catalog No.	Size
Pinpoint™ Slide RNA Isolation System I	R1003	50 preps.

Individual Kit Components	Catalog No.	Amount
Pinpoint™ Solution	D3001-1	1 ml
RNA Extraction Buffer	R1003-2-12 R1003-2-50	12 ml 50 ml
RNA Wash Buffer (concentrate)	R1003-2-6 R1003-3-12	6 ml 12 ml
DNase/RNase-Free Water	W1001-6 W1001-10	6 ml 10 ml
Zymo-Spin™ IC Columns	C1004-50	50
Collection Tubes	C1001-50	50

Complete Your Workflow

- ✓ For tough-to-lyse samples, use ZR BashingBead Lysis Tubes:

ZR BashingBead Lysis Tubes	
2.0 mm beads #S6003	Plant/animal tissue
0.1 + 0.5 mm beads #S6012	Microbes
0.1 + 2.0 mm beads #S6014	Microbes in tissue/insects

- ✓ For isolation of RNA from any sample:

Quick-RNA kits	
Miniprep Plus #R1057/R1058	$\leq 10^7$ cells, ≤ 50 mg tissue
MagBeads #R2132/R2133	Automatable (Tecan, Hamilton, Kingfisher, etc.)

- ✓ For clean-up (purification) and concentration of any RNA sample. (e.g., from the aqueous phase of TRIzol[®] extractions) or from any enzymatic reaction (e.g., DNase I treated RNA):

RNA Clean & Concentrator kits	
Microprep #R1013-R1014	DNase I Set included
MagBeads #R1082	Automatable (Tecan, Hamilton, Kingfisher, etc.)

- ✓ For NGS:

Zymo-Seq RiboFree Total RNA Library Prep kit	
#R3000	12 preps
#R3003	96 preps

Troubleshooting Guide

Problem	Possible Causes and Suggested Solutions
RNA degradation	RNA is very susceptible to RNase digestion; thus we encourage the use of freshly prepared tissue sections. If a sample cannot be processed immediately, store it at $\leq -70^{\circ}\text{C}$ or submerge it in a 95% ethanol bath at -20°C . Processing of tissue sections stored for ≥ 1 month at room temperature is not recommended. If the eluted RNA will not be used immediately it is recommended that 1 U/10 μl of RNase inhibitor be added to the sample prior to storage at -70°C .
Insufficient RNA	Make sure an appropriate sampling area is selected for processing. Select an area of the tissue that will contain ≥ 50 cells. Increase the sampling area if the tissue type contains few cells (e.g., fatty tissue and connective tissue). The sampling size can vary from 1 mm^2 to over 100 mm^2 . We recommend that the sample thickness be $\geq 10 \mu\text{m}$.
RT/qPCR parameters are not optimized	It is recommended that the conditions used for RT/qPCR be optimized prior to using template RNA purified by the Pinpoint™ Slide RNA Isolation System I . It may be necessary to increase both the annealing and extension times and adjust the number of cycles for low copy number mRNAs.
DNA contamination	To remove DNA: - Perform DNase I treatment post-purification. See RNA Clean & Concentrator kit #R1013.

For technical assistance, please contact 1-888-882-9682 or email tech@zymoresearch.com

Notes



**100% satisfaction guarantee on all Zymo Research products,
or your money back.**

Zymo Research is committed to simplifying your research with quality products and services. If you are dissatisfied with this product for any reason, please call 1(888) 882-9682.

Integrity of kit components is guaranteed for up to one year from date of purchase. Reagents are routinely tested on a lot-to-lot basis to ensure they provide the highest performance and reliability.

This product is for research use only and should only be used by trained professionals. It is not for use in diagnostic procedures. Some reagents included with this kit are irritants. Wear protective gloves and eye protection. Follow the safety guidelines and rules enacted by your research institution or facility.

™ Trademarks of Zymo Research Corporation
Pinpoint® is a registered trademark of Zymo Research Corporation.



ZYMO RESEARCH

The **BEAUTY** of **SCIENCE** is to Make Things **SIMPLE**®



tech@zymoresearch.com



www.zymoresearch.com



Toll Free: (888) 882-9682