



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



ZYMO RESEARCH

RNA
Purification
Made Simple

Quick-DNA/RNA™ Pathogen Miniprep

DNA & RNA from any vector-borne pathogen

Highlights

- Spin-column purification of pathogen (virus, bacteria, protozoa) DNA and RNA from a wide variety of vectors (mosquitoes, fleas, ticks, etc.) and tissue types (mammals, birds, etc.).
- High-quality DNA/RNA is ready for Next-Gen sequencing, RT/qPCR, hybridization, etc.
- DNA/RNA Shield™ is included for sample collection, inactivation, storage and preservation.

Catalog Numbers:
R1042, R1043



Scan with your smart-phone camera to
view the online protocol/video.



tech@zymoresearch.com



www.zymoresearch.com



Toll Free: (888) 882-9682

Table of Contents

Product Contents	01
Specifications	02
Product Description	03
Workflow	04
Protocol	05
(I) Buffer Preparation	05
(II) Sample Preparation.....	06
Tissue, Swabs, DNA/RNA Shield Samples	06
(III) DNA/RNA Purification.....	07
Appendices	08
DNase I Treatment	08
Ordering Information	09
Complete Your Workflow	10
Troubleshooting Guide	11
Notes	12
Guarantee	13

Product Contents

Quick-DNA/RNA™ Pathogen Miniprep	R1042 (50 prep)	R1043 (200 prep)
DNA/RNA Shield™	50 ml	250 ml
Pathogen DNA/RNA Buffer ¹	50 ml	100 ml (x2)
Proteinase K ² (lyophilized) w/ Storage Buffer	5 mg	20 mg
Zymo-Spin™ IICR Columns	50	200
Collection Tubes	50	200
Pathogen DNA/RNA Wash Buffer ³ (concentrate)	6 ml (x2)	48 ml
ZymoBIOMICS™ DNase/RNase-Free Water	3 ml	10 ml (x2)
Instruction Manual	1 pc	1 pc
ZR Bashing Bead™ Lysis Tubes (sold separately)	S6014-50 (50 pack)	

Storage Temperature - Store all kit components (i.e., buffers, columns) at room temperature.

Before use:

1 Add beta-mercaptoethanol to 0.5% (v/v) *i.e.*, add 250 µl or 500 µl β-Me per 50 ml or 100 ml **Pathogen DNA/RNA Buffer**.

2 Reconstitute lyophilized **Proteinase K** according to page 4, Buffer Preparation. Store frozen aliquots.

3 Add 24 ml of 100% ethanol (26 ml of 95% ethanol) to the 6 ml **Pathogen DNA/RNA Wash Buffer** concentrate (R1042) or 192 ml of 100% ethanol (204 ml of 95% ethanol) to the 48 ml **Pathogen DNA/RNA Wash Buffer** concentrate (R1043).

Specifications

- **Sample Sources** – ≤ 10 mg vectors (mosquitoes, fleas, ticks, other tough-to-lyse insects) and tissue types (animal tissue, plants, other hosts) or up to 400 μ l liquid sample (e.g., samples in DNA/RNA Shield™).

For samples in UTM®/VTM®, PBS or saline, see Sample Preparation, page 6.

- **Purity** – High-quality DNA and RNA is ready for Next-Gen Sequencing, RT/qPCR, etc.
- **Binding Capacity** – 50 μ g DNA/RNA **Zymo-Spin™ IICR Columns**.
- **Elution Volume** – ≥ 25 μ l **ZymoBIOMICS™ DNase/RNase-Free Water**.
- **Equipment Needed** (user provided) – Beta-mercaptoethanol (b-Me), Ethanol (95-100%), Microcentrifuge.
- **Materials** (available separately) –

ZR BashingBead™ Lysis Tubes (S6014-50; 0.1 & 2.0 mm beads)
DNA/RNA Shield™ (R1200; 2X concentrate)
DNase I Set (E1010; 50 rxns.; 250 U DNase I (lyophilized)
supplied w/ DNA Digestion Buffer, 4 ml)
DNA/RNA Prep Buffer (D7010-2-50; 50 ml)


Product Description

The **Quick-DNA/RNA™ Pathogen Miniprep** kit is a spin-column purification of pathogen (virus, bacteria, protozoa) DNA and RNA from a wide variety of vectors (mosquitoes, fleas, ticks, etc.) and tissue types (mammals, birds, etc.) collected, transported and stored in **DNA/RNA Shield™**. DNA/RNA Shield™ is used for nucleic acid preservation and inactivation of pathogens.

The kit features a storage/lysis buffer system and can be combined with high density ZR BashingBead™ Lysis Tubes (*recommended) to facilitate complete homogenization of hard-to-lyse samples for efficient nucleic acid isolation. Small (>50 nt) and large (> 200 kb) DNA and RNA are bound to the column, washed and then eluted.

The isolated high-quality nucleic acids are suitable for all downstream applications such as Next-Gen sequencing, hybridization-based and RT/qPCR detection.

Inhibitor-free Detection of Viruses in Mosquitoes



West Nile Virus (top) and La Crosse Encephalitis (bottom) viral particles were spiked-in mosquito homogenate, purified using the **Quick-DNA/RNA™ Pathogen** kit and then detected by RT-qPCR.

Workflow

Pathogen

virus, bacteria, protozoa

Vectors

insects, mammals, birds, etc.

Biological samples

biopsies, fecal, blood, etc.



DNA/RNA Shield™

add to sample for collection, transport and storage



*recommended:
Bashing Bead Lysis

Spin-column

catalog no. R1042, R1043

MagBead

catalog no. R2145, R2146

Bind, Wash and Elute!



DNA/RNA is ready for

NGS, (RT)PCR, etc.

Protocol

The protocol consists of: (I) Buffer Preparation, (II) Sample Preparation and (III) DNA/RNA Purification.

(I) Buffer Preparation

- ✓ Add beta-mercaptoethanol to 0.5% (v/v) i.e., add 250 μ l or 500 μ l β -Me per 50 ml or 100 ml **Pathogen DNA/RNA Buffer**, respectively.
- ✓ Add 24 ml of 100% ethanol (26 ml of 95% ethanol) to the 6 ml **Pathogen DNA/RNA Wash Buffer** concentrate (R1042) or 192 ml of 100% ethanol (204 ml of 95% ethanol) to the 48 ml **Pathogen DNA/RNA Wash Buffer** concentrate (R1043).
- ✓ Add 260 μ l or 1,040 μ l **Proteinase K Storage Buffer** to reconstitute per 5 mg or 20 mg lyophilized **Proteinase K**, respectively. Mix by vortexing. Use immediately or store frozen aliquots.

(II) Sample Preparation

- ✓ Perform all steps at room temperature (20-30°C).
- ✓ Up to 400 µl sample can be processed per prep.

Tissue¹ (tough-to-lyse insects and tissue)

Up to 10 mg tissue (see table below) can be homogenized in ≥ 400 µl **DNA/RNA Shield**^{TM2} with a mortar/pestle, dounce, syringe, tissue grinder, or bead beating with a high-speed homogenizer. To remove particulate debris, centrifuge the homogenate and transfer the cleared supernatant into a nuclease-free tube. Proceed with purification, page 7.

Recommended: To a ZR BashingBeadTM Lysis Tube (#S6014; sold separately), add the appropriate amount of sample to **DNA/RNA Shield**TM, then mechanically homogenize³.

Input	Mosquitoes	Ticks	Fleas	Deer fly	Tissue
per prep (pool)	50	1 engorged 5 flat adults 20 nymphs	10	1 adult	animal, plant (≤ 10 mg)
High-speed ³	3-5 minutes				30-60 seconds
Low-speed ³	≥ 10 minutes (optimization may be required)				

Swabs (UTM[®]/VTM[®], PBS, saline, etc.)

Proceed directly with purification, page 7.

Optional: To inactivate, store and preserve samples at room temperature prior to processing, add an equal volume of DNA/RNA ShieldTM (2X concentrate) (#R1200; sold separately) to a volume of liquid sample (1:1) and mix well.

Samples in DNA/RNA Shield^{TM2} **collection devices** (swabs, saliva, etc.)

Proceed directly with purification, page 7.

Optional - **Proteinase K treatment** (protein-rich samples e.g., tissue and biological liquids, can be treated):

Add 1% **Proteinase K** (v/v) at 20 mg/ml directly to a liquid sample. Mix well and incubate at room temperature for 15 minutes. Note: Up to 5% Proteinase K can be added (e.g., tissue). For example: Add 4-20 µl Proteinase K to each 400 µl sample.

1 To remove particulate debris or cryoprecipitates (if any), centrifuge and transfer up to 400 µl of the cleared supernatant into a nuclease-free plate/tube (not provided).

2 At this point, samples in DNA/RNA ShieldTM can be stored at ambient temperature (4-25°C) for a month, 3 days at 37°C, or long-term (> 1 year) -20°C or below.

3 Perform bead beating at maximum speed with a high-speed homogenizer (e.g., MP Bio FastPrep-24, Bertin Precellys, etc.) or with a low-speed homogenizer (e.g., Vortex Genie, etc.).

(III) DNA/RNA Purification

- ✓ Perform all steps at room temperature (20-30°C).
 - ✓ The sample input can be scaled up or down, proportionally.
1. Add 800 µl **Pathogen DNA/RNA Buffer**¹ to each 400 µl sample² (2:1) and mix well².
 2. Transfer the mixture into a **Zymo-Spin™ IICR Column**³ in a **Collection Tube** and centrifuge. Discard the flow-through.

Optional: At this point, DNase I treatment can be performed (see Appendices, page 8).
 3. Add 500 µl **Pathogen DNA/RNA Wash Buffer** to the column and centrifuge. Discard the flow-through and repeat this step.
 4. Add 500 µl ethanol (95-100%) to the column and centrifuge for 1 minute to ensure removal of any residual ethanol. Discard the collection tube and carefully transfer the column into a new nuclease-free tube (not provided).
 5. To elute DNA/RNA, add 50 µl **ZymoBIOMICS™ DNase/RNase-Free Water** directly to the matrix of the column and centrifuge.

Alternatively, for highly concentrated DNA/RNA use ≥ 25 µl elution volume.

The eluted DNA/RNA⁴ can be used immediately or stored frozen.

1 To ensure efficient lysis and deproteinization, up to 5 volumes of Pathogen DNA/RNA Buffer can be used per volume of sample.

2 Up to 400 µl sample can be processed per prep.

3 For volumes > 700 µl, column can be reloaded.

4 It is recommended to titrate the DNA/RNA eluate for downstream applications (i.e., RT/qPCR, etc.).

Appendices

DNase I Treatment

- ✓ For DNA-free RNA, DNase I treatment can be performed using DNase I Set (E1010; 50 reactions), DNA/RNA Prep Buffer (D7010-2-50) and DNA/RNA Wash Buffer (concentrate) (D7010-3-6); materials sold separately.

For each sample to be treated, prepare **DNase I Reaction Mix** in an RNase-free tube (not provided) and mix by gentle inversion:

DNase I Reaction Mix	
DNA Digestion Buffer	75 µl
DNase I (reconstituted; 1 U/ul) ^{1,2}	5 µl

1. Following DNA/RNA binding (page 7, step 2), add 400 µl **DNA/RNA Wash Buffer**³ to the column, centrifuge and discard the flow-through.
2. Add 80 µl **DNase I Reaction Mix** directly to the matrix of the column.
3. Incubate at room temperature for (20-30°C) for 15 minutes.
4. Add 500 µl **DNA/RNA Prep Buffer** to the column, centrifuge and discard the flow-through.
5. Proceed with DNA/RNA Purification (page 7, step 3).

1 Prior to use, reconstitute lyophilized 250 U **DNase I** (E1009-A) to 1U/µl (final concentration) with 275 µl nuclease-free water (not provided), mix by gentle inversion and store frozen aliquots.

2 Unit definition – one unit increases the absorbance of a high molecular weight DNA solution at a rate of 0.001 A₂₆₀ units/ml of reaction mixture at 25°C.

3 Before use, add 24 ml of 100% ethanol (26 ml of 95% ethanol) to the 6 ml **DNA/RNA Wash Buffer** concentrate.

Ordering Information

Product Description	Catalog No.	Size
Quick-DNA/RNA™ Pathogen Miniprep	R1042 R1043	50 preps. 200 preps.

Individual Kit Components	Catalog No.	Amount
Pathogen DNA/RNA Buffer	R1042-1-50 R1042-1-100	50 ml 100 ml
Proteinase K Set supplied w/ Storage Buffer	D3001-2-5 D3001-2-20	5 mg 20 mg
Zymo-Spin™ IICR Columns	C1078-50 C1078-250	50 250
Collection Tubes (2 ml)	C1001-50 C1001-500	50 500
Pathogen DNA/RNA Wash Buffer (concentrate)	R1042-2-6 R1042-2-48	6 ml 48 ml
ZymoBIOMICS™ DNase/RNase-Free Water	D4302-5-3 D4302-5-10	3 ml 10 ml
DNA/RNA Shield™	R1100-50 R1100-250	50 ml 250 ml
DNA/RNA Shield™ Fecal Collection Tube	R1101	10
DNA/RNA Shield™ Collection Tube	R1102	50
DNA/RNA Shield™ Lysis Tube (microbe)	R1103	50
DNA/RNA Shield™ Lysis Tube (microbe) w/ swab	R1104	50
DNA/RNA Shield™ Lysis Tube (tissue)	R1105	50
DNA/RNA Shield™ Collection Tube w/ Swab (1 ml fill)	R1106 R1107	10 50
DNA/RNA Shield™ Collection Tube w/ Swab (2 ml fill)	R1108 R1109	10 50
DNA/RNA Shield™ Saliva Collection Kit (2 ml fill)	R1210	1
DNase I Set (250 U DNase I (lyophilized) supplied with DNA Digestion Buffer, 4 ml)	E1010	1
DNA/RNA Prep Buffer	D7010-2-50	50 ml

Complete Your Workflow

- ✓ For sample collection, inactivation of pathogens, storage and preservation of nucleic acids, use DNA/RNA Shield™ collection devices:

DNA/RNA Shield™ Collection Devices	
DNA/RNA Shield™ Collection Tube w/ Swab (1 ml fill or 2 ml fill) #R1107, R1109	For swab samples of nasal, throat, etc.
DNA/RNA Shield™ Saliva Collection Kit (2 ml fill) #R1210	For saliva, sputum, etc.
DNA/RNA Shield™ Collection Tube DNA/RNA Shield™ Lysis Tube (microbe) DNA/RNA Shield™ Lysis Tube (microbe) w/ swab DNA/RNA Shield™ Lysis Tube (tissue) #R1102-R1105	For microbes, tissue, etc. (2 ml lysis tubes used for bead beating homogenization)

- ✓ For RNA clean-up (purification) from the aqueous phase (e.g., TRIzol, TRI Reagent or similar) or from any enzymatic reaction (e.g., DNase I treated RNA):

RNA Clean & Concentrator	
Microprep #R1013, R1015	DNase I Set included (#R1013)
MagBeads #R1081, R1082	(#R1082)

Troubleshooting Guide

Problem	Possible Causes and Suggested Solutions
RNA degradation	<p>To prevent RNA degradation:</p> <p>Immediately collect and lyse fresh sample into a stabilization reagent (i.e., DNA/RNA Shield™) to ensure nucleic acid stability. Homogenized samples in DNA/RNA Shield™ can be stored frozen for later processing.</p>
Low nucleic acid content and/or low sensitivity in downstream application	<p>Incomplete deproteinization due to high-protein content in the sample (tissue, biological liquids, etc.):</p> <ul style="list-style-type: none">- Increase the volume of DNA/RNA Shield™ added to the sample.- Perform Proteinase K treatment (see Sample Preparation, page 5). <p>Increase eluate input:</p> <ul style="list-style-type: none">-Titrate the DNA/RNA eluate for downstream applications (i.e., RT/qPCR).
DNA contamination	<p>To remove DNA:</p> <ul style="list-style-type: none">- Perform DNase I treatment during the purification (page 8) or perform DNase I treatment post-purification (#R1017), then clean-up the treated sample.

For technical assistance, please contact 1-888-882-9682 or email tech@zymoresearch.com



**100% satisfaction guarantee on all Zymo Research products,
or your money back.**

Zymo Research is committed to simplifying your research with quality products and services. If you are dissatisfied with this product for any reason, please call 1(888) 882-9682.

Integrity of kit components is guaranteed for up to one year from date of purchase. Reagents are routinely tested on a lot-to-lot basis to ensure they provide the highest performance and reliability.

This product is for research use only and should only be used by trained professionals. It is not for use in diagnostic procedures. Some reagents included with this kit are irritants. Wear protective gloves and eye protection. Follow the safety guidelines and rules enacted by your research institution or facility.

TM Trademarks of Zymo Research Corporation
Quick-DNA/RNA[®] is a registered trademark of Zymo Research Corporation. Other trademarks: Tecan[®], Hamilton[®], Thermo Fisher[®]



ZYMO RESEARCH

The **BEAUTY** of **SCIENCE** is to Make Things **SIMPLE**®



tech@zymoresearch.com



www.zymoresearch.com



Toll Free: (888) 882-9682