



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



ZYMO RESEARCH

DNA
Purification
MADE SIMPLE

Frozen-EZ Yeast Transformation II™

Fast procedure to make competent yeast cells with high transformation efficiency.

Highlights

- High transformation efficiency that yields approximately 10^5 - 10^6 transformants per μg plasmid DNA (circular).
- Frozen storage of competent cells for future use.
- Fast and easy steps to make competent yeast cells within 10 minutes.
- Easy, single step transformation procedure that takes less than an hour.
- Broad spectrum (*S. cerevisiae*, *C. albicans*, *S. pombe*, *Pichia pastoris*).
- Simple method for multiple plasmid transformations.

Catalog Numbers:
T2001



Scan with your smart-phone camera to
view the online protocol/video.



tech@zymoresearch.com



www.zymoresearch.com



Toll Free: (888) 882-9682

Table of Contents

Product Contents	01
Product Description	02
Standard Transformation Protocol	03
Preparation of Competent Cells	03
Transformation	03
Micro-Scale Protocol for Multiple-Plasmid Transformations	04
Preparation of Competent Cells	04
Transformation	04
Appendix	05
FAQ's	07
Ordering Information	09

Product Contents

Frozen-EZ Yeast Transformation II™	T2001 (120 Rxns.)	Storage
Frozen-EZ Yeast Solution 1	60 ml	0 – 4°C
Frozen-EZ Yeast Solution 2	6 ml	0 – 4°C
Frozen-EZ Yeast Solution 3	60 ml	0 – 4°C
Instruction Manual	1	-

Reagents in this kit are designed for 120 regular or 600 micro-scale transformation experiments.

Product Description

The **Frozen-EZ Yeast Transformation II™** is designed to make yeast transformations easier and more efficient than protocols that are currently in wide usage. Version II further improves the efficiency and reliability of its predecessor. The competent yeast cells prepared with these reagents can be used immediately for transformation experiments or can be stored frozen at or below -70°C for direct use in the future. This method is suitable for both circular and linear plasmid transformations.

Protocol

Standard Transformation Protocol

Note: For maximum transformation efficiency, please read "Optimizing Your Conditions for Higher Efficiency," pg. 5. If your experiment requires larger scale than the "Standard Transformation Protocol" you can increase the volume in each step proportionally according to this procedure.

Preparation of Competent Cells

Grow yeast cells at 30°C in 10 ml YPD broth until mid-log phase ($\sim 5 \times 10^6$ - 2×10^7 cells/ml or OD₆₀₀ of 0.8-1.0). The following steps are accomplished at room temperature.

1. Pellet the cells at 500 x g for 4 minutes and discard the supernatant.
2. Add 10 ml **Frozen-EZ Yeast Solution 1** to wash the pellet. Repellet the cells and discard the supernatant.
3. Add 1 ml **Frozen-EZ Yeast Solution 2** to resuspend the pellet.

At this point, the competent cells can be used for transformations directly or stored frozen at or below -70°C for future use. It is important to freeze the cells slowly. To accomplish this, either wrap the aliquotted cells in 2-6 layers of paper towels or place in a Styrofoam box before placing in the freezer. **DO NOT** use liquid nitrogen to snap-freeze the cells.

Transformation

This part of the procedure is the same for both frozen stored (thawed at room temperature) and freshly prepared competent yeast cells.

1. Mix 50 μ l of competent cells with 0.2-1 μ g DNA (in less than 5 μ l volume); add 500 μ l **Frozen-EZ Yeast Solution 3** and mix thoroughly.
2. Incubate at 30°C for 45 minutes. Mix vigorously by flicking with finger or vortexing (if appropriate for your DNA) 2-3 times during this incubation.

3. Spread 50-150 μl of the above transformation mixture on an appropriate plate. It is unnecessary to pellet and wash the cells before spreading.
4. Incubate the plates at 30°C for 2-4 days to allow for growth of transformants.

Attention: For transformations of *C. albicans*, use freshly prepared competent cells; frozen cells sometimes give poor results.

Micro-Scale Protocol for Multiple-Plasmid Transformations

Note: For maximum transformation efficiency, please read "Optimizing Your Conditions for Higher Efficiency" on page 5.

This protocol can generate enough transformants for most applications. If your work requires a large number of transformants as in library screening, use "Standard Transformation Protocol" (page 3) and "Optimizing Your Conditions for Higher Efficiency" on page 5.

Preparation of Competent Cells: Same as "Standard Transformation Protocol" (see page 3).

Transformation: This can be done either in 96-well plates or tubes.

1. Add 10 μl competent cells to DNA (0.2-1 μg), mix by tapping or lightly vortexing; add 100 μl **Frozen-EZ Yeast Solution 3** and mix thoroughly.
2. Incubate at 30°C for 60-90 minutes. Mix vigorously 2-3 times during this incubation.
3. Directly spread the transformation mixture onto 1-2 plates.
4. Incubate the plates at 30°C for 2-4 days to allow for growth of transformants.

Appendix

Optimizing Your Conditions for Higher Efficiency

Cell growth conditions, strain differences, and other factors may influence the transformation efficiency. The following factors need to be considered if your experiments require high transformation efficiency:

1. Cell Growth State

Cells used should be in mid-log phase. Early or late log-phase cells yield relatively fewer transformants.

2. Cell Density

Optimal cell density is between 5×10^6 and 2×10^7 cells/ml (0.8-1.0 OD₆₀₀), but in most cases use of cultures with cell densities at the high end of the range greatly increases transformation efficiency in comparison to cultures with cell densities at the low end of the range. The optimal cell density can also be achieved by resuspending the cells in a smaller volume of **Frozen-EZ Yeast Solution 2** (for example, harvesting 10 ml cultures in low-density range and resuspending in 0.5 ml of **Frozen-EZ Yeast Solution 2** instead of 1 ml as in the "Standard Transformation Protocol").

3. Incubation Time After Adding Frozen-EZ Yeast Solution 3

Although an incubation time of 45 minutes is good for general purposes, transformation efficiency is much better with longer incubation times (up to 2-3 hours in most cases). Again, results vary according to strain used.

4. Medium Used for Plating

Not all commercially available media are created equal. The difference in transformation efficiency using different grade of media can be several folds. Our test results show that media from Difco are the most reliable.

5. DNA Amount Used

For transformation experiments using circular DNA such as 2 μ -based plasmids, we find that the efficiency stops increasing linearly as you increase the DNA above 1 μ g using the standard transformation protocol. For integrative transformation, purity and amount of DNA used are important. Higher amount of linearized DNA is recommended to achieve the best results (up to 5 μ g of DNA can be used in the "Standard Transformation Protocol").

6. Pre-Warm Solutions

Usually, it is not necessary to pre-warm all the solutions. For most purposes, cold solutions from the refrigerator will work fine. But, if your experiment requires maximum transformation efficiency, such as in library screening, the pre-warming of the solutions before the transformation experiment will result in higher efficiency. Calculate and measure the amounts of each solution that you will need to complete your experiment. Pre-warm these amounts to 20-37°C before doing experiments.

7. Outgrowth After Transformation

Some selection markers and certain yeast strains need outgrowth step in rich medium for high efficiency transformation after Step 3 of Transformation, page 3. For this, add 4 volumes of YPD medium to the transformation mixture at the end of transformation, (2 ml YPD for the Standard Transformation, page 3) mix, incubate at 30°C for 2 hours. After this incubation, spin the cells down, resuspend the cells in 2 ml of water, spin the cells down again to wash the cells, resuspend the cells in original (0.5 ml for “Standard Transformation Protocol”) or less volume of water for plating on selection plates.

FAQ's

1. **What is in this kit and how does it work?**

The procedure utilized in this kit is designed, in some ways, similar to the lithium cation based method. No spheroplast step is involved. The mechanism probably involves some metabolic pathways that we do not fully understand.

2. **Does this kit work on *C. albicans*, *S. pombe*, or *Pichia pastoris*?**

Yes. Based on data from other labs and ours, this kit does work as well on *C. albicans*, *S. pombe*, or *Pichia pastoris* as well as *S. cerevisiae*.

3. **How long can I store my competent yeast cells below -70°C?**

There is no loss of transformation efficiency after half a year of storage below -70°C. The transformation efficiency gradually starts to drop after 6 months.

4. **How should I thaw the stored frozen competent cells?**

Thaw at room temperature.

5. **Do the frequencies of freezing and thawing affect the transformation efficiency?**

We usually see a 10-30% increase of transformants after the first cycle of freezing. You can refreeze and thaw the competent yeast cells 3-4 times without noticeable effect on the transformation efficiency. Further freezing-thawing cycles adversely affect the transformation efficiency.

6. **In step 2 of the transformation, do I need to incubate strictly at 30°C?**

No. Temperatures between 30°C and 37°C are in the optimal temperature range. Incubation below or above this range greatly reduces the transformation efficiency.

7. **Can I use DNA directly from restriction enzyme digestions without purification?**

Yes. Different digestion buffers have only a slight effect on the transformation efficiency. You should try to keep the DNA volume in 5 µl per transformation experiment by increasing the concentration of DNA in the digestion reaction.

8. How much transformation mixture should I spread on a selective plate?

For most circular plasmid transformations, 50 μ l is enough. But if you use linearized DNA or use more than one selection marker, you can apply up to 200 μ l of transformation mixture on each plate to increase the number of transformants. The number of transformants increases linearly with the amount of transformation mixture applied to each plate.

9. Do I need to use plates with sorbitol?

No. Use any plate that is appropriate for your experiment.

10. Do I need to add carrier DNA?

No. There is no need for carrier DNA.

Note: Most of the data above is based on the testing of *S. cerevisiae*.

Ordering Information

Product Description	Catalog No.	Size
Frozen-EZ Yeast Transformation II™	T2001	120 Rxns.

Individual Kit Components	Catalog No.	Amount
Frozen-EZ Yeast Solution I	T2002	60 ml
Frozen-EZ Yeast Solution II	T2003	6 ml
Frozen-EZ Yeast Solution III	T2004	60 ml



**100% satisfaction guarantee on all Zymo Research products,
or your money back.**

Zymo Research is committed to simplifying your research with quality products and services. If you are dissatisfied with this product for any reason, please call 1(888) 882-9682.

Integrity of kit components is guaranteed for up to one year at 4°C from date of purchase. Reagents are routinely tested on a lot-to-lot basis to ensure they provide the highest performance and reliability.

This product is for research use only and should only be used by trained professionals. It is not for use in diagnostic procedures. Some reagents included with this kit are irritants. Wear protective gloves and eye protection. Follow the safety guidelines and rules enacted by your research institution or facility.

™ Trademarks of Zymo Research Corporation



ZYMO RESEARCH

The **BEAUTY** of **SCIENCE** is to Make Things **SIMPLE**®



tech@zymoresearch.com



www.zymoresearch.com



Toll Free: (888) 882-9682